

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 February 2002 (07.02.2002)

PCT

(10) International Publication Number
WO 02/10387 A2

(51) International Patent Classification⁷: **C12N 15/12**,
C07K 14/705, 16/28, C12Q 1/68, G01N 33/50, 33/577,
A61K 38/17, A01K 67/027

(21) International Application Number: PCT/US01/23433

(22) International Filing Date: 25 July 2001 (25.07.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/221,478 27 July 2000 (27.07.2000) US
60/223,268 3 August 2000 (03.08.2000) US
60/227,054 21 August 2000 (21.08.2000) US
60/231,121 8 September 2000 (08.09.2000) US
60/232,243 13 September 2000 (13.09.2000) US
60/232,691 15 September 2000 (15.09.2000) US
60/235,146 22 September 2000 (22.09.2000) US

(71) Applicant (for all designated States except US): **INCYTE GENOMICS, INC.** [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **THORNTON, Michael** [US/US]; 9 Medway Road, Woodside, CA 94062 (US). **PATTERSON, Chandra** [US/US]; 490 Sherwood Way, #1, Menlo Park, CA 94025 (US). **LAL, Preeti** [IN/US]; P.O. Box 5142, Santa Clara, CA 95056 (US). **BURFORD, Neil** [GB/US]; 105 Wildwood Circle, Durham, CT 06422 (US). **YUE, Henry** [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). **GANDHI, Ameena, R.** [US/US]; 837 Roble Avenue, #1, Redwood City, CA 94025 (US). **ELLIOT, Vicki, S.** [US/US]; 3770 Polton Place Way, San Jose, CA 95121 (US). **RAMKUMAR, Jayalaxmi** [IN/US]; 34359 Maybird Circle, Fremont, CA 94555 (US). **BAUGHN, Mariah, R.** [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). **KALLICK, Deborah, A.** [US/US]; 900 Olive Street, Menlo Park, CA 94025 (US). **WALIA, Narinder, K.** [US/US]; 890 Davis

Street #205, San Leandro, CA 94577 (US). **HAFALIA, April, J., A.** [US/US]; 2227 Calle de Primavera, Santa Clara, CA 95054 (US). **YAO, Monique, G.** [US/US]; 111 Frederick Court, Mountain View, CA 94043 (US). **LU, Yan** [CN/US]; 3885 Corrina Way, Palo Alto, CA 94303 (US). **TRIBOULEY, Catherine, M.** [FR/US]; 1121 Tennessee Street, #5, San Francisco, CA 94107 (US). **POLICKY, Jennifer, L.** [US/US]; 1511 Jarvis Court, San Jose, CA 95118 (US). **KEARNEY, Liam** [IE/US]; 50 Woodside Avenue, San Francisco, CA 94127 (US). **GRAUL, Richard, C.** [—/US]; 682-29th Avenue, San Francisco, CA 94121 (US). **WARREN, Bridget, A.** [US/US]; 10130 Parwood Drive #2, Cupertino, CA 95014 (US). **LEE, Ernestine, A.** [US/US]; 624 Kains Street, Albany, CA 94706 (US). **DING, Li** [US/US]; 3353 Alma Street, #146, Palo Alto, CA 94306 (US).

(74) Agents: **HAMLET-COX, Diana et al.**; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: G-PROTEIN COUPLED RECEPTORS

(57) Abstract: The invention provides human G-protein coupled receptors (GCREC) and polynucleotides which identify and encode GCREC. The invention also provides expression vectors, host cells, antibodies, agonist, and antagonist. The invention also provides methods for diagnosing, treating, or preventing disorders associated with aberrant expression of GCREC.

WO 02/10387 A2

G-PROTEIN COUPLED RECEPTORS

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of G-protein coupled
5 receptors and to the use of these sequences in the diagnosis, treatment, and prevention of cell
proliferative, neurological, cardiovascular, gastrointestinal, autoimmune/inflammatory, and metabolic
disorders, and viral infections, and in the assessment of the effects of exogenous compounds on the
expression of nucleic acid and amino acid sequences of G-protein coupled receptors.

BACKGROUND OF THE INVENTION

Signal transduction is the general process by which cells respond to extracellular signals.
Signal transduction across the plasma membrane begins with the binding of a signal molecule, e.g., a
hormone, neurotransmitter, or growth factor, to a cell membrane-receptor. The receptor, thus
activated, triggers an intracellular biochemical cascade that ends with the activation of an intracellular
15 target molecule, such as a transcription factor. This process of signal transduction regulates all types
of cell functions including cell proliferation, differentiation, and gene transcription. The G-protein
coupled receptors (GPCRs), encoded by one of the largest families of genes yet identified, play a
central role in the transduction of extracellular signals across the plasma membrane. GPCRs have a
proven history of being successful therapeutic targets.

20 GPCRs are integral membrane proteins characterized by the presence of seven hydrophobic
transmembrane domains which together form a bundle of antiparallel alpha (α) helices. GPCRs range
in size from under 400 to over 1000 amino acids (Strosberg, A.D. (1991) Eur. J. Biochem. 196:1-10;
Coughlin, S.R. (1994) Curr. Opin. Cell Biol. 6:191-197). The amino-terminus of a GPCR is
extracellular, is of variable length, and is often glycosylated. The carboxy-terminus is cytoplasmic
25 and generally phosphorylated. Extracellular loops alternate with intracellular loops and link the
transmembrane domains. Cysteine disulfide bridges linking the second and third extracellular loops
may interact with agonists and antagonists. The most conserved domains of GPCRs are the
transmembrane domains and the first two cytoplasmic loops. The transmembrane domains account,
in part, for structural and functional features of the receptor. In most cases, the bundle of α helices
30 forms a ligand-binding pocket. The extracellular N-terminal segment, or one or more of the three
extracellular loops, may also participate in ligand binding. Ligand binding activates the receptor by
inducing a conformational change in intracellular portions of the receptor. In turn, the large, third
intracellular loop of the activated receptor interacts with a heterotrimeric guanine nucleotide binding
(G) protein complex which mediates further intracellular signaling activities, including the activation
35 of second messengers such as cyclic AMP (cAMP), phospholipase C, and inositol triphosphate, and

the interaction of the activated GPCR with ion channel proteins. (See, e.g., Watson, S. and S. Arkinstall (1994) The G-protein Linked Receptor Facts Book, Academic Press, San Diego CA, pp. 2-6; Bolander, F.F. (1994) Molecular Endocrinology, Academic Press, San Diego CA, pp. 162-176; Baldwin, J.M. (1994) *Curr. Opin. Cell Biol.* 6:180-190.)

5 GPCRs include receptors for sensory signal mediators (e.g., light and olfactory stimulatory molecules); adenosine, γ -aminobutyric acid (GABA), hepatocyte growth factor, melanocortins, neuropeptide Y, opioid peptides, opsins, somatostatin, tachykinins, vasoactive intestinal polypeptide family, and vasopressin; biogenic amines (e.g., dopamine, epinephrine and norepinephrine, histamine, glutamate (metabotropic effect), acetylcholine (muscarinic effect), and serotonin); chemokines; lipid
10 mediators of inflammation (e.g., prostaglandins and prostanoids, platelet activating factor, and leukotrienes); and peptide hormones (e.g., bombesin, bradykinin, calcitonin, C5a anaphylatoxin, endothelin, follicle-stimulating hormone (FSH), gonadotropic-releasing hormone (GnRH), neurokinin, and thyrotropin-releasing hormone (TRH), and oxytocin). GPCRs which act as receptors for stimuli that have yet to be identified are known as orphan receptors.

15 The diversity of the GPCR family is further increased by alternative splicing. Many GPCR genes contain introns, and there are currently over 30 such receptors for which splice variants have been identified. The largest number of variations are at the protein C-terminus. N-terminal and cytoplasmic loop variants are also frequent, while variants in the extracellular loops or transmembrane domains are less common. Some receptors have more than one site at which variance
20 can occur. The splicing variants appear to be functionally distinct, based upon observed differences in distribution, signaling, coupling, regulation, and ligand binding profiles (Kilpatrick, G.J. et al. (1999) *Trends Pharmacol. Sci.* 20:294-301).

GPCRs can be divided into three major subfamilies: the rhodopsin-like, secretin-like, and metabotropic glutamate receptor subfamilies. Members of these GPCR subfamilies share similar
25 functions and the characteristic seven transmembrane structure, but have divergent amino acid sequences. The largest family consists of the rhodopsin-like GPCRs, which transmit diverse extracellular signals including hormones, neurotransmitters, and light. Rhodopsin is a photosensitive GPCR found in animal retinas. In vertebrates, rhodopsin molecules are embedded in membranous stacks found in photoreceptor (rod) cells. Each rhodopsin molecule responds to a photon of light by
30 triggering a decrease in cGMP levels which leads to the closure of plasma membrane sodium channels. In this manner, a visual signal is converted to a neural impulse. Other rhodopsin-like GPCRs are directly involved in responding to neurotransmitters. These GPCRs include the receptors for adrenaline (adrenergic receptors), acetylcholine (muscarinic receptors), adenosine, galanin, and glutamate (N-methyl-D-aspartate/NMDA receptors). (Reviewed in Watson, S. and S. Arkinstall
35 (1994) The G-Protein Linked Receptor Facts Book, Academic Press, San Diego CA, pp. 7-9, 19-22,

32-35, 130-131, 214-216, 221-222; Habert-Ortoli, E. et al. (1994) Proc. Natl. Acad. Sci. USA 91:9780-9783.)

The galanin receptors mediate the activity of the neuroendocrine peptide galanin, which inhibits secretion of insulin, acetylcholine, serotonin and noradrenaline, and stimulates prolactin and growth hormone release. Galanin receptors are involved in feeding disorders, pain, depression, and Alzheimer's disease (Kask, K. et al. (1997) Life Sci. 60:1523-1533). Other nervous system rhodopsin-like GPCRs include a growing family of receptors for lysophosphatidic acid and other lysophospholipids, which appear to have roles in development and neuropathology (Chun, J. et al. (1999) Cell Biochem. Biophys. 30:213-242).

10 The largest subfamily of GPCRs, the olfactory receptors, are also members of the rhodopsin-like GPCR family. These receptors function by transducing odorant signals. Numerous distinct olfactory receptors are required to distinguish different odors. Each olfactory sensory neuron expresses only one type of olfactory receptor, and distinct spatial zones of neurons expressing distinct receptors are found in nasal passages. For example, the RA1c receptor which was isolated from a rat brain library, has been shown to be limited in expression to very distinct regions of the brain and a defined zone of the olfactory epithelium (Raming, K. et al. (1998) Receptors Channels 6:141-151). However, the expression of olfactory-like receptors is not confined to olfactory tissues. For example, three rat genes encoding olfactory-like receptors having typical GPCR characteristics showed expression patterns not only in taste and olfactory tissue, but also in male reproductive tissue
15 (Thomas, M.B. et al. (1996) Gene 178:1-5).

Members of the secretin-like GPCR subfamily have as their ligands peptide hormones such as secretin, calcitonin, glucagon, growth hormone-releasing hormone, parathyroid hormone, and vasoactive intestinal peptide. For example, the secretin receptor responds to secretin, a peptide hormone that stimulates the secretion of enzymes and ions in the pancreas and small intestine
25 (Watson, supra, pp. 278-283). Secretin receptors are about 450 amino acids in length and are found in the plasma membrane of gastrointestinal cells. Binding of secretin to its receptor stimulates the production of cAMP.

Examples of secretin-like GPCRs implicated in inflammation and the immune response include the EGF module-containing, mucin-like hormone receptor (Emr1) and CD97 receptor
30 proteins. These GPCRs are members of the recently characterized EGF-TM7 receptors subfamily. These seven transmembrane hormone receptors exist as heterodimers in vivo and contain between three and seven potential calcium-binding EGF-like motifs. CD97 is predominantly expressed in leukocytes and is markedly upregulated on activated B and T cells (McKnight, A.J. and S. Gordon (1998) J. Leukoc. Biol. 63:271-280).

35 The third GPCR subfamily is the metabotropic glutamate receptor family. Glutamate is the

major excitatory neurotransmitter in the central nervous system. The metabotropic glutamate receptors modulate the activity of intracellular effectors, and are involved in long-term potentiation (Watson, supra, p.130). The Ca^{2+} -sensing receptor, which senses changes in the extracellular concentration of calcium ions, has a large extracellular domain including clusters of acidic amino acids which may be involved in calcium binding. The metabotropic glutamate receptor family also includes pheromone receptors, the GABA_B receptors, and the taste receptors.

Other subfamilies of GPCRs include two groups of chemoreceptor genes found in the nematodes Caenorhabditis elegans and Caenorhabditis briggsae, which are distantly related to the mammalian olfactory receptor genes. The yeast pheromone receptors STE2 and STE3, involved in the response to mating factors on the cell membrane, have their own seven-transmembrane signature, as do the cAMP receptors from the slime mold Dictyostelium discoideum, which are thought to regulate the aggregation of individual cells and control the expression of numerous developmentally-regulated genes.

GPCR mutations, which may cause loss of function or constitutive activation, have been associated with numerous human diseases (Coughlin, supra). For instance, retinitis pigmentosa may arise from mutations in the rhodopsin gene. Furthermore, somatic activating mutations in the thyrotropin receptor have been reported to cause hyperfunctioning thyroid adenomas, suggesting that certain GPCRs susceptible to constitutive activation may behave as protooncogenes (Parma, J. et al. (1993) Nature 365:649-651). GPCR receptors for the following ligands also contain mutations associated with human disease: lutenizing hormone (precocious puberty); vasopressin V_2 (X-linked nephrogenic diabetes); glucagon (diabetes and hypertension); calcium (hyperparathyroidism, hypocalcuria, hypercalcemia); parathyroid hormone (short limbed dwarfism); β_3 -adrenoceptor (obesity, non-insulin-dependent diabetes mellitus); growth hormone releasing hormone (dwarfism); and adrenocorticotropin (glucocorticoid deficiency) (Wilson, S. et al. (1998) Br. J. Pharmacol. 125:1387-1392; Stadel, J.M. et al. (1997) Trends Pharmacol. Sci. 18:430-437). GPCRs are also involved in depression, schizophrenia, sleeplessness, hypertension, anxiety, stress, renal failure, and several cardiovascular disorders (Horn, F. and G. Vriend (1998) J. Mol. Med. 76:464-468).

In addition, within the past 20 years several hundred new drugs have been recognized that are directed towards activating or inhibiting GPCRs. The therapeutic targets of these drugs span a wide range of diseases and disorders, including cardiovascular, gastrointestinal, and central nervous system disorders as well as cancer, osteoporosis and endometriosis (Wilson, supra; Stadel, supra). For example, the dopamine agonist L-dopa is used to treat Parkinson's disease, while a dopamine antagonist is used to treat schizophrenia and the early stages of Huntington's disease. Agonists and antagonists of adrenoceptors have been used for the treatment of asthma, high blood pressure, other cardiovascular disorders, and anxiety; muscarinic agonists are used in the treatment of glaucoma and

tachycardia; serotonin 5HT_{1D} antagonists are used against migraine; and histamine H₁ antagonists are used against allergic and anaphylactic reactions, hay fever, itching, and motion sickness (Horn, supra).

Recent research suggests potential future therapeutic uses for GPCRs in the treatment of metabolic disorders including diabetes, obesity, and osteoporosis. For example, mutant V2 vasopressin receptors causing nephrogenic diabetes could be functionally rescued in vitro by co-expression of a C-terminal V2 receptor peptide spanning the region containing the mutations. This result suggests a possible novel strategy for disease treatment (Schöneberg, T. et al. (1996) EMBO J. 15:1283-1291). Mutations in melanocortin-4 receptor (MC4R) are implicated in human weight regulation and obesity. As with the vasopressin V2 receptor mutants, these MC4R mutants are defective in trafficking to the plasma membrane (Ho, G. and R.G. MacKenzie (1999) J. Biol. Chem. 274:35816-35822), and thus might be treated with a similar strategy. The type 1 receptor for parathyroid hormone (PTH) is a GPCR that mediates the PTH-dependent regulation of calcium homeostasis in the bloodstream. Study of PTH/receptor interactions may enable the development of novel PTH receptor ligands for the treatment of osteoporosis (Mannstadt, M. et al. (1999) Am. J. Physiol. 277:F665-F675).

The chemokine receptor group of GPCRs have potential therapeutic utility in inflammation and infectious disease. (For review, see Locati, M. and P.M. Murphy (1999) Annu. Rev. Med. 50:425-440.) Chemokines are small polypeptides that act as intracellular signals in the regulation of leukocyte trafficking, hematopoiesis, and angiogenesis. Targeted disruption of various chemokine receptors in mice indicates that these receptors play roles in pathologic inflammation and in autoimmune disorders such as multiple sclerosis. Chemokine receptors are also exploited by infectious agents, including herpesviruses and the human immunodeficiency virus (HIV-1) to facilitate infection. A truncated version of chemokine receptor CCR5, which acts as a coreceptor for infection of T-cells by HIV-1, results in resistance to AIDS, suggesting that CCR5 antagonists could be useful in preventing the development of AIDS.

The discovery of new G-protein coupled receptors, and the polynucleotides encoding them, satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of cell proliferative, neurological, cardiovascular, gastrointestinal, autoimmune/inflammatory, and metabolic disorders, and viral infections, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of G-protein coupled receptors.

SUMMARY OF THE INVENTION

The invention features purified polypeptides, G-protein coupled receptors, referred to

collectively as "GCREC" and individually as "GCREC-1," "GCREC-2," "GCREC-3," "GCREC-4," "GCREC-5," "GCREC-6," "GCREC-7," "GCREC-8," "GCREC-9," "GCREC-10," "GCREC-11," "GCREC-12," "GCREC-13," "GCREC-14," "GCREC-15," "GCREC-16," "GCREC-17," "GCREC-18," and "GCREC-19." In one aspect, the invention provides an isolated polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-19.

The invention further provides an isolated polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19. In one alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-19. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:20-38.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, b) a polypeptide comprising a naturally occurring amino acid sequence at least

90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19. The method comprises a) 5 culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid 10 sequence selected from the group consisting of SEQ ID NO:1-19, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group 15 consisting of SEQ ID NO:1-19.

The invention further provides an isolated polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:20-38, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID 20 NO:20-38, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide selected from the group 25 consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:20-38, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:20-38, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The 30 method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex, and optionally, if 35 present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous

nucleotides.

The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of
5 SEQ ID NO:20-38, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:20-38, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain
10 reaction amplification, and b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

The invention further provides a composition comprising an effective amount of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, b) a polypeptide comprising a
15 naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, and a pharmaceutically acceptable excipient. In one embodiment, the
20 composition comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-19. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional GCREC, comprising administering to a patient in need of such treatment the composition.

The invention also provides a method for screening a compound for effectiveness as an
25 agonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, and d) an
30 immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of
35 treating a disease or condition associated with decreased expression of functional GCREC,

comprising administering to a patient in need of such treatment the composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, b) a polypeptide
5 comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19. The method comprises a) exposing a sample comprising the
10 polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional GCRC, comprising administering to a patient in need of such treatment the composition.

15 The invention further provides a method of screening for a compound that specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, c) a biologically active fragment of a polypeptide
20 having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19. The method comprises a) combining the polypeptide with at least one test compound under suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

25 The invention further provides a method of screening for a compound that modulates the activity of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, c) a biologically active fragment of a polypeptide
30 having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19. The method comprises a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the

polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

The invention further provides a method for screening a compound for effectiveness in
 5 altering expression of a target polynucleotide, wherein said target polynucleotide comprises a polynucleotide sequence selected from the group consisting of SEQ ID NO:20-38, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

The invention further provides a method for assessing toxicity of a test compound, said
 10 method comprising a) treating a biological sample containing nucleic acids with the test compound; b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:20-38, ii) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a
 15 polynucleotide sequence selected from the group consisting of SEQ ID NO:20-38, iii) a polynucleotide having a sequence complementary to i), iv) a polynucleotide complementary to the polynucleotide of ii), and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide selected from the group consisting of i) a
 20 polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:20-38, ii) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:20-38, iii) a polynucleotide complementary to the polynucleotide of i), iv) a polynucleotide complementary to the polynucleotide of ii), and v) an RNA equivalent of i)-iv). Alternatively, the target
 25 polynucleotide comprises a fragment of a polynucleotide sequence selected from the group consisting of i)-v) above; c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

30

BRIEF DESCRIPTION OF THE TABLES

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the present invention.

Table 2 shows the GenBank identification number and annotation of the nearest GenBank homolog for polypeptides of the invention. The probability score for the match between each polypeptide and its GenBank homolog is also shown.

Table 3 shows structural features of polypeptide sequences of the invention, including
5 predicted motifs and domains, along with the methods, algorithms, and searchable databases used for analysis of the polypeptides.

Table 4 lists the cDNA and/or genomic DNA fragments which were used to assemble polynucleotide sequences of the invention, along with selected fragments of the polynucleotide sequences.

10 Table 5 shows the representative cDNA library for polynucleotides of the invention.

Table 6 provides an appendix which describes the tissues and vectors used for construction of the cDNA libraries shown in Table 5.

Table 7 shows the tools, programs, and algorithms used to analyze the polynucleotides and polypeptides of the invention, along with applicable descriptions, references, and threshold
15 parameters.

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these
20 may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a
25 reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs.
30 Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the
35 invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

"GCREC" refers to the amino acid sequences of substantially purified GCREC obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

5 The term "agonist" refers to a molecule which intensifies or mimics the biological activity of GCREC. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of GCREC either by directly interacting with GCREC or by acting on components of the biological pathway in which GCREC participates.

10 An "allelic variant" is an alternative form of the gene encoding GCREC. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times
15 in a given sequence.

"Altered" nucleic acid sequences encoding GCREC include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as GCREC or a polypeptide with at least one functional characteristic of GCREC. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide
20 probe of the polynucleotide encoding GCREC, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding GCREC. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent GCREC. Deliberate amino acid substitutions may be made on the basis of similarity in
25 polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of GCREC is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine.
30 Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to a sequence of a naturally occurring
35 protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid

sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

5 The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of GCREC. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of GCREC either by directly interacting with GCREC or by acting on components of the biological pathway in which GCREC participates.

10 The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind GCREC polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the
15 translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

 The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that
20 makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

25 The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having
30 modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the
35 designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic GCREC, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

"Complementary" describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement, 3'-TCA-5'.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding GCREC or fragments of GCREC may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (Applied Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
30	Ala	Gly, Ser
	Arg	His, Lys
	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
35	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
	Gly	Ala

	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
	Leu	Ile, Val
	Lys	Arg, Gln, Glu
5	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
	Thr	Ser, Val
	Trp	Phe, Tyr
10	Tyr	His, Phe, Trp
	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to a chemically modified polynucleotide or polypeptide. Chemical modifications of a polynucleotide can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

"Differential expression" refers to increased or upregulated; or decreased, downregulated, or absent gene or protein expression, determined by comparing at least two different samples. Such comparisons may be carried out between, for example, a treated and an untreated sample, or a diseased and a normal sample.

"Exon shuffling" refers to the recombination of different coding regions (exons). Since an exon may represent a structural or functional domain of the encoded protein, new proteins may be assembled through the novel reassortment of stable substructures, thus allowing acceleration of the evolution of new protein functions.

A "fragment" is a unique portion of GCREC or the polynucleotide encoding GCREC which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A

fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids
5 selected from the first 250 or 500 amino acids (or first 25% or 50%) of a polypeptide as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:20-38 comprises a region of unique polynucleotide sequence that
10 specifically identifies SEQ ID NO:20-38, for example, as distinct from any other sequence in the genome from which the fragment was obtained. A fragment of SEQ ID NO:20-38 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:20-38 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:20-38 and the region of SEQ ID NO:20-38 to which the fragment corresponds are routinely
15 determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-19 is encoded by a fragment of SEQ ID NO:20-38. A fragment of SEQ ID NO:1-19 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-19. For example, a fragment of SEQ ID NO:1-19 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-19. The precise length of
20 a fragment of SEQ ID NO:1-19 and the region of SEQ ID NO:1-19 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A "full length" polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A
25 "full length" polynucleotide sequence encodes a "full length" polypeptide sequence.

"Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a
30 standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e
35 sequence alignment program. This program is part of the LASERGENE software package, a suite of

molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequences.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

20 *Matrix: BLOSUM62*
 Reward for match: 1
 Penalty for mismatch: -2
 Open Gap: 5 and Extension Gap: 2 penalties
 Gap x drop-off: 50
 25 *Expect: 10*
 Word Size: 11
 Filter: on

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

35 Nucleic acid sequences that do not show a high degree of identity may nevertheless encode

similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Open Gap: 11 and Extension Gap: 1 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 3

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size and which contain all of the elements required for

chromosome replication, segregation and maintenance.

The term "humanized antibody" refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

5 "Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the
10 stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity.
15 Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic
20 strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

25 High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents
30 include, for instance, sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such
35 similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C_0t or R_0t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

An "immunogenic fragment" is a polypeptide or oligopeptide fragment of GCREC which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of GCREC which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of GCREC. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of GCREC.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of

amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

“Post-translational modification” of an GCREC may involve lipidation, glycosylation, phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by cell type depending on the enzymatic milieu of GCREC.

“Probe” refers to nucleic acid sequences encoding GCREC, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes. “Primers” are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al. (1987) Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al. (1990) PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of

Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, *supra*. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription, translation, or RNA stability.

"Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent, chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and

other moieties known in the art.

An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing GCREC, nucleic acids encoding GCREC, or fragments thereof may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acid residues or nucleotides by different amino acid residues or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

A "transcript image" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

"Transformation" describes a process by which exogenous DNA is introduced into a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed cells" includes stably transformed cells in which the inserted DNA is capable of

replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook et al. (1989), supra.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides will generally have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of

the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence

5 identity over a certain defined length of one of the polypeptides.

THE INVENTION

The invention is based on the discovery of new human G-protein coupled receptors (GCREC), the polynucleotides encoding GCREC, and the use of these compositions for the diagnosis, 10 treatment, or prevention of cell proliferative, neurological, cardiovascular, gastrointestinal, autoimmune/inflammatory, and metabolic disorders, and viral infections.

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the invention. Each polynucleotide and its corresponding polypeptide are correlated to a single Incyte project identification number (Incyte Project ID). Each polypeptide sequence is denoted 15 by both a polypeptide sequence identification number (Polypeptide SEQ ID NO:) and an Incyte polypeptide sequence number (Incyte Polypeptide ID) as shown. Each polynucleotide sequence is denoted by both a polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and an Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) as shown.

Table 2 shows sequences with homology to the polypeptides of the invention as identified by 20 BLAST analysis against the GenBank protein (genpept) database. Columns 1 and 2 show the polypeptide sequence identification number (Polypeptide SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for polypeptides of the invention. Column 3 shows the GenBank identification number (Genbank ID NO:) of the nearest GenBank homolog. Column 4 shows the probability score for the match between each polypeptide and its GenBank 25 homolog. Column 5 shows the annotation of the GenBank homolog.

Table 3 shows various structural features of the polypeptides of the invention. Columns 1 and 2 show the polypeptide sequence identification number (SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for each polypeptide of the invention. Column 3 shows the number of amino acid residues in each polypeptide. Column 4 shows potential 30 phosphorylation sites, and column 5 shows potential glycosylation sites, as determined by the MOTIFS program of the GCG sequence analysis software package (Genetics Computer Group, Madison WI). Column 6 shows amino acid residues comprising signature sequences, domains, and motifs. Column 7 shows analytical methods for protein structure/function analysis and in some cases, searchable databases to which the analytical methods were applied.

35 Together, Tables 2 and 3 summarize the properties of each polypeptide of the invention, and

these properties establish that the claimed polypeptides are G-protein coupled receptors. For example, SEQ ID NO:1 is 40% identical to Meleagris gallopavo G protein-coupled P2Y nucleotide receptor (GenBank ID g2707256) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 4.0e-62, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:1 also contains a rhodopsin family 7 transmembrane receptor domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS and BLAST analyses provide further corroborative evidence that SEQ ID NO:1 is G-protein coupled receptor. SEQ ID NO:2 was analyzed and annotated in a similar manner. These analyses indicate that SEQ ID NO:2 is a pheromone receptor (Dulac, C. and R. Axel (1995) Cell 83:195-206).

As a further example, SEQ ID NO:6 is 29% identical to human C-C chemokine receptor type 1 (GenBank ID g179985) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 1.6e-15, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:6 also contains a 7 transmembrane receptor (rhodopsin family) domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO:6 is a chemokine receptor.

As a further example, SEQ ID NO:9 is 95% identical to rat calcium-independent alpha-latrotoxin receptor (GenBank ID g3882981) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 0.0, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:9 also contains a 7-transmembrane receptor (secretin family) domain and a latrophilin/CL-1-like GPS domain, as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS, MOTIFS, and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO:9 is a latrophilin-related G-protein coupled receptor.

As a further example, SEQ ID NO:12 is 84% identical to Mus musculus G-protein coupled receptor GPR73 (GenBank ID g7248884) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 6.7e-166, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:12 also contains a 7 transmembrane receptor (rhodopsin family) domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS analysis reveals the presence of a rhodopsin-like

GPCR superfamily signature (See Table 3). Additional data from MOTIFS and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO:12 is a G-protein coupled receptor.

As a further example, SEQ ID NO:15 is 80% identical to rat serotonin receptor (GenBank ID g310075) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is $2.5e-152$, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:15 also contains a rhodopsin family receptor domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS, analyses provide further corroborative evidence that SEQ ID NO:15 is a G-protein coupled receptor.

As a further example, SEQ ID NO:16 is 71% identical to mouse olfactory receptor E3 (GenBank ID g3983382) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is $1.9e-88$, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:16 also contains a rhodopsin family 7-transmembrane receptor domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS, MOTIFS, and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO:16 is an olfactory G-protein coupled receptor.

As a further example, SEQ ID NO:17 is 83% identical to mouse olfactory G-protein coupled receptor G3 (GenBank ID g3983398) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is $5.0e-99$, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:17 also contains a rhodopsin family 7-transmembrane receptor domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS, MOTIFS, and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO:17 is an olfactory G-protein coupled receptor. SEQ ID NO:2-5, SEQ ID NO:7-8, SEQ ID NO:10-11, SEQ ID NO:13-14, and SEQ ID NO:18-19 were analyzed and annotated in a similar manner. The algorithms and parameters for the analysis of SEQ ID NO:1-19 are described in Table 7.

As shown in Table 4, the full length polynucleotide sequences of the present invention were assembled using cDNA sequences or coding (exon) sequences derived from genomic DNA, or any combination of these two types of sequences. Columns 1 and 2 list the polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and the corresponding Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) for each polynucleotide of the invention. Column 3 shows the length of each polynucleotide sequence in basepairs. Column 4 lists fragments

of the polynucleotide sequences which are useful, for example, in hybridization or amplification technologies that identify SEQ ID NO:20-38 or that distinguish between SEQ ID NO:20-38 and related polynucleotide sequences. Column 5 shows identification numbers corresponding to cDNA sequences, coding sequences (exons) predicted from genomic DNA, and/or sequence assemblages comprised of both cDNA and genomic DNA. These sequences were used to assemble the full length polynucleotide sequences of the invention. Columns 6 and 7 of Table 4 show the nucleotide start (5') and stop (3') positions of the cDNA and/or genomic sequences in column 5 relative to their respective full length sequences.

The identification numbers in Column 5 of Table 4 may refer specifically, for example, to Incyte cDNAs along with their corresponding cDNA libraries. For example, 7075196H1 is the identification number of an Incyte cDNA sequence, and BRAUTDR04 is the cDNA library from which it is derived. Incyte cDNAs for which cDNA libraries are not indicated were derived from pooled cDNA libraries (e.g., 71906055V1). Alternatively, the identification numbers in column 5 may refer to GenBank cDNAs or ESTs (e.g., g900324) which contributed to the assembly of the full length polynucleotide sequences. In addition, the identification numbers in column 5 may identify sequences derived from the ENSEMBL (The Sanger Centre, Cambridge, UK) database (*i.e.*, those sequences including the designation "ENST"). Alternatively, the identification numbers in column 5 may be derived from the NCBI RefSeq Nucleotide Sequence Records Database (*i.e.*, those sequences including the designation "NM" or "NT") or the NCBI RefSeq Protein Sequence Records (*i.e.*, those sequences including the designation "NP"). Alternatively, the identification numbers in column 5 may refer to assemblages of both cDNA and Genscan-predicted exons brought together by an "exon stitching" algorithm. For example, FL_XXXXXX_N₁_N₂_YYYYY_N₃_N₄ represents a "stitched" sequence in which XXXXXX is the identification number of the cluster of sequences to which the algorithm was applied, and YYYYY is the number of the prediction generated by the algorithm, and N_{1,2,3...}, if present, represent specific exons that may have been manually edited during analysis (See Example V). Alternatively, the identification numbers in column 5 may refer to assemblages of exons brought together by an "exon-stretching" algorithm. For example, FLXXXXXX_gAAAAA_gBBBBB_1_N is the identification number of a "stretched" sequence, with XXXXXX being the Incyte project identification number, gAAAAA being the GenBank identification number of the human genomic sequence to which the "exon-stretching" algorithm was applied, gBBBBB being the GenBank identification number or NCBI RefSeq identification number of the nearest GenBank protein homolog, and N referring to specific exons (See Example V). In instances where a RefSeq sequence was used as a protein homolog for the "exon-stretching" algorithm, a RefSeq identifier (denoted by "NM," "NP," or "NT") may be used in place of the GenBank identifier (*i.e.*, gBBBBB).

Alternatively, a prefix identifies component sequences that were hand-edited, predicted from genomic DNA sequences, or derived from a combination of sequence analysis methods. The following Table lists examples of component sequence prefixes and corresponding sequence analysis methods associated with the prefixes (see Example IV and Example V).

5	Prefix	Type of analysis and/or examples of programs
	GNN, GFG, ENST	Exon prediction from genomic sequences using, for example, GENSCAN (Stanford University, CA, USA) or FGENES (Computer Genomics Group, The Sanger Centre, Cambridge, UK).
	GBI	Hand-edited analysis of genomic sequences.
	FL	Stitched or stretched genomic sequences (see Example V).
10	INCY	Full length transcript and exon prediction from mapping of EST sequences to the genome. Genomic location and EST composition data are combined to predict the exons and resulting transcript.

In some cases, Incyte cDNA coverage redundant with the sequence coverage shown in column 5 was obtained to confirm the final consensus polynucleotide sequence, but the relevant Incyte cDNA identification numbers are not shown.

15 Table 5 shows the representative cDNA libraries for those full length polynucleotide sequences which were assembled using Incyte cDNA sequences. The representative cDNA library is the Incyte cDNA library which is most frequently represented by the Incyte cDNA sequences which were used to assemble and confirm the above polynucleotide sequences. The tissues and vectors which were used to construct the cDNA libraries shown in Table 5 are described in Table 6.

20 The invention also encompasses GCREC variants. A preferred GCREC variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the GCREC amino acid sequence, and which contains at least one functional or structural characteristic of GCREC.

25 The invention also encompasses polynucleotides which encode GCREC. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:20-38, which encodes GCREC. The polynucleotide sequences of SEQ ID NO:20-38, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

30 The invention also encompasses a variant of a polynucleotide sequence encoding GCREC. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at

least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding GCREC. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:20-38 which has at least about 70%, or alternatively at least about 85%, or even at least about 5 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:20-38. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of GCREC.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding GCREC, some bearing minimal 10 similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring GCREC, and all such variations are to be considered 15 as being specifically disclosed.

Although nucleotide sequences which encode GCREC and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring GCREC under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding GCREC or its derivatives possessing a substantially different codon usage, e.g., inclusion of non- 20 naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding GCREC and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a 25 greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode GCREC and GCREC derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to 30 introduce mutations into a sequence encoding GCREC or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:20-38 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 35 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in

"Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Applied Biosystems), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Applied Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Applied Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding GCREC may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) *Nucleic Acids Res.* 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) *PCR Methods Applic.* 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) *Nucleic Acids Res.* 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 primer analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of

about 68°C to 72°C.

When screening for full length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Applied Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode GCREC may be cloned in recombinant DNA molecules that direct expression of GCREC, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express GCREC.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter GCREC-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Cramer, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or improve the biological properties of GCREC, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired

properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, sequences encoding GCREC may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) *Nucleic Acids Symp. Ser. 7*:215-223; and Horn, T. et al. (1980) *Nucleic Acids Symp. Ser. 7*:225-232.) Alternatively, GCREC itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Applied Biosystems). Additionally, the amino acid sequence of GCREC, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, supra, pp. 28-53.)

In order to express a biologically active GCREC, the nucleotide sequences encoding GCREC or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding GCREC. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding GCREC. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding GCREC and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-

frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

5 Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding GCREC and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding GCREC. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast
15 transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. (See, e.g., Sambrook, supra; Ausubel, supra; Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509; Engelhard, E.K. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3224-3227; Sandig, V. et al. (1996) *Hum. Gene Ther.* 7:1937-1945; Takamatsu, N. (1987) *EMBO J.* 6:307-311; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659; and Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355.) Expression vectors derived
25 from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. (See, e.g., Di Nicola, M. et al. (1998) *Cancer Gen. Ther.* 5(6):350-356; Yu, M. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90(13):6340-6344; Buller, R.M. et al. (1985) *Nature* 317(6040):813-815; McGregor, D.P. et al. (1994) *Mol. Immunol.* 31(3):219-226; and Verma, I.M. and N. Somia (1997) *Nature* 389:239-242.) The invention is not limited by the host cell employed.

30 In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding GCREC. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding GCREC can be achieved using a multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding GCREC into the vector's multiple
35 cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of

transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of GCREC are needed, e.g. for the production of antibodies, vectors which direct high level expression of GCREC may be used. For example, vectors
5 containing the strong, inducible SP6 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of GCREC. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such
10 vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, supra; Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; and Scorer, C.A. et al. (1994) Bio/Technology 12:181-184.)

Plant systems may also be used for expression of GCREC. Transcription of sequences encoding GCREC may be driven by viral promoters, e.g., the 35S and 19S promoters of CaMV used
15 alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105.)
20 These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding GCREC may be ligated into
25 an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses GCREC in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-
30 based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet.
35 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of GCREC in cell lines is preferred. For example, sequences encoding GCREC can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector.

- 5 Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.
- 10 Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk*⁻ and *apv*⁻ cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to
- 15 methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc.
- 20 Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

- 25 Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding GCREC is inserted within a marker gene sequence, transformed cells containing sequences encoding GCREC can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding GCREC under the control of a
- 30 single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

- In general, host cells that contain the nucleic acid sequence encoding GCREC and that express GCREC may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR
- 35 amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or

chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of GCREC using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and
5 fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on GCREC is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and
10 Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding GCREC
15 include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding GCREC, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety
20 of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding GCREC may be cultured under
25 conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode GCREC may be designed to contain signal sequences which direct secretion of GCREC through a prokaryotic or eukaryotic cell membrane.

30 In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity.
35 Different host cells which have specific cellular machinery and characteristic mechanisms for

post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding GCREC may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric GCREC protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of GCREC activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the GCREC encoding sequence and the heterologous protein sequence, so that GCREC may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, *supra*, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled GCREC may be achieved *in vitro* using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

GCREC of the present invention or fragments thereof may be used to screen for compounds that specifically bind to GCREC. At least one and up to a plurality of test compounds may be screened for specific binding to GCREC. Examples of test compounds include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

In one embodiment, the compound thus identified is closely related to the natural ligand of GCREC, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, e.g., Coligan, J.E. et al. (1991) Current Protocols in Immunology 1(2): Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which GCREC binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the

compound can be rationally designed using known techniques. In one embodiment, screening for these compounds involves producing appropriate cells which express GCREC, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, Drosophila, or E. coli. Cells expressing GCREC or cell membrane fractions which contain GCREC are then
5 contacted with a test compound and binding, stimulation, or inhibition of activity of either GCREC or the compound is analyzed.

An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with GCREC, either in
10 solution or affixed to a solid support, and detecting the binding of GCREC to the compound. Alternatively, the assay may detect or measure binding of a test compound in the presence of a labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

GCREC of the present invention or fragments thereof may be used to screen for compounds that modulate the activity of GCREC. Such compounds may include agonists, antagonists, or partial or inverse agonists. In one embodiment, an assay is performed under conditions permissive for GCREC activity, wherein GCREC is combined with at least one test compound, and the activity of GCREC in the presence of a test compound is compared with the activity of GCREC in the absence
20 of the test compound. A change in the activity of GCREC in the presence of the test compound is indicative of a compound that modulates the activity of GCREC. Alternatively, a test compound is combined with an in vitro or cell-free system comprising GCREC under conditions suitable for GCREC activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of GCREC may do so indirectly and need not come in direct contact with the
25 test compound. At least one and up to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding GCREC or their mammalian homologs may be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent Number 5,175,383 and U.S. Patent Number
30 5,767,337.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (neo; Capecchi, M.R. (1989) Science 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes
35 place using the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-

specific manner (Marth, J.D. (1996) Clin. Invest. 97:1999-2002; Wagner, K.U. et al. (1997) Nucleic Acids Res. 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce
5 heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

Polynucleotides encoding GCREC may also be manipulated in vitro in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate
10 into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) Science 282:1145-1147).

Polynucleotides encoding GCREC can also be used to create "knockin" humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding GCREC is injected into animal ES cells, and the injected
15 sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress GCREC, e.g., by secreting GCREC in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) Biotechnol. Annu. Rev. 4:55-
20 74).

THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of GCREC and G-protein coupled receptors. In addition, the expression of GCREC is closely associated with brain tissue, fetal brain tissue, colon polyps, diseased colon tissue, colon
25 tumor tissue, diseased gallbladder tissue, heart tissue, diseased breast tissue, interleukin-5 stimulated eosinophils, tumor tissue, and reproductive tissues. Therefore, GCREC appears to play a role in cell proliferative, neurological, cardiovascular, gastrointestinal, autoimmune/inflammatory, and metabolic disorders, and viral infections. In the treatment of disorders associated with increased GCREC expression or activity, it is desirable to decrease the expression or activity of GCREC. In the
30 treatment of disorders associated with decreased GCREC expression or activity, it is desirable to increase the expression or activity of GCREC.

Therefore, in one embodiment, GCREC or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GCREC. Examples of such disorders include, but are not limited to, a cell proliferative
35 disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed

connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, colon, gall bladder, ganglia, 5 gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, 10 retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, 15 tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, 20 myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; a cardiovascular disorder such as arteriovenous fistula, atherosclerosis, hypertension, vasculitis, 25 Raynaud's disease, aneurysms, arterial dissections, varicose veins, thrombophlebitis and phlebothrombosis, vascular tumors, complications of thrombolysis, balloon angioplasty, vascular replacement, and coronary artery bypass graft surgery, congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve, mitral annular 30 calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis, nonbacterial thrombotic endocarditis, endocarditis of systemic lupus erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart disease, congenital heart disease, and complications of cardiac transplantation; a gastrointestinal disorder such as dysphagia, peptic esophagitis, esophageal spasm, esophageal stricture, esophageal carcinoma,

- dyspepsia, indigestion, gastritis, gastric carcinoma, anorexia, nausea, emesis, gastroparesis, antral or pyloric edema, abdominal angina, pyrosis, gastroenteritis, intestinal obstruction, infections of the intestinal tract, peptic ulcer, cholelithiasis, cholecystitis, cholestasis, pancreatitis, pancreatic carcinoma, biliary tract disease, hepatitis, hyperbilirubinemia, cirrhosis, passive congestion of the liver, hepatoma, infectious colitis, ulcerative colitis, ulcerative proctitis, Crohn's disease, Whipple's disease, Mallory-Weiss syndrome, colonic carcinoma, colonic obstruction, irritable bowel syndrome, short bowel syndrome, diarrhea, constipation, gastrointestinal hemorrhage, acquired immunodeficiency syndrome (AIDS) enteropathy, jaundice, hepatic encephalopathy, hepatorenal syndrome, hepatic steatosis, hemochromatosis, Wilson's disease, α_1 -antitrypsin deficiency, Reye's syndrome, primary sclerosing cholangitis, liver infarction, portal vein obstruction and thrombosis, centrilobular necrosis, peliosis hepatis, hepatic vein thrombosis, veno-occlusive disease, preeclampsia, eclampsia, acute fatty liver of pregnancy, intrahepatic cholestasis of pregnancy, and hepatic tumors including nodular hyperplasias, adenomas, and carcinomas; an autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a metabolic disorder such as diabetes, obesity, and osteoporosis; and an infection by a viral agent classified as adenovirus, arenavirus, bunyavirus, calicivirus, coronavirus, filovirus, hepadnavirus, herpesvirus, flavivirus, orthomyxovirus, parvovirus, papovavirus, paramyxovirus, picornavirus, poxvirus, reovirus, retrovirus, rhabdovirus, and togavirus.

In another embodiment, a vector capable of expressing GCREC or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GCREC including, but not limited to, those described above.

In a further embodiment, a composition comprising a substantially purified GCREC in

conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GCREC including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of GCREC may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GCREC including, but not limited to, those listed above.

In a further embodiment, an antagonist of GCREC may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of GCREC. Examples of such disorders include, but are not limited to, those cell proliferative, neurological, cardiovascular, gastrointestinal, autoimmune/inflammatory, and metabolic disorders, and viral infections described above. In one aspect, an antibody which specifically binds GCREC may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express GCREC.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding GCREC may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of GCREC including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of GCREC may be produced using methods which are generally known in the art. In particular, purified GCREC may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind GCREC. Antibodies to GCREC may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with GCREC or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic

polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to GCREC have an amino acid sequence consisting of at least about 5 amino acids, and generally will
5 consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of GCREC amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to GCREC may be prepared using any technique which provides for
10 the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120.)

15 In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda, S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single
20 chain antibodies may be adapted, using methods known in the art, to produce GCREC-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte
25 population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. USA 86:3833-3837; Winter, G. et al. (1991) Nature 349:293-299.)

Antibody fragments which contain specific binding sites for GCREC may also be generated. For example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin
30 digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) Science 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired
35 specificity. Numerous protocols for competitive binding or immunoradiometric assays using either

polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between GCREC and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering GCREC epitopes is generally used, but a competitive binding assay
5 may also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for GCREC. Affinity is expressed as an association constant, K_a , which is defined as the molar concentration of GCREC-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions.
10 The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple GCREC epitopes, represents the average affinity, or avidity, of the antibodies for GCREC. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular GCREC epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12} L/mole are preferred for use in immunoassays in
15 which the GCREC-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of GCREC, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).
20

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation
25 of GCREC-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding GCREC, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications
30 of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA, RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding GCREC. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding GCREC. (See, e.g., Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press Inc.,
35 Totawa NJ.)

In therapeutic use, any gene delivery system suitable for introduction of the antisense sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g.,

5 Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995) 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, *supra*; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other

10 systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.* 25(14):2730-2736.)

In another embodiment of the invention, polynucleotides encoding GCREC may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency

15 (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaese, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene*

20 *Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, familial hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal, R.G. (1995) *Science* 270:404-410; Verma, I.M. and N. Somia (1997) *Nature* 389:239-242)), (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated cell proliferation), or (iii) express a protein which affords protection against intracellular parasites

25 (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) *Nature* 335:395-396; Poeschla, E. et al. (1996) *Proc. Natl. Acad. Sci. USA.* 93:11395-11399), hepatitis B or C virus (HBV, HCV); fungal parasites, such as Candida albicans and Paracoccidioides brasiliensis; and protozoan parasites such as Plasmodium falciparum and Trypanosoma cruzi). In the case where a genetic deficiency in GCREC expression or regulation causes disease, the expression of

30 GCREC from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in GCREC are treated by constructing mammalian expression vectors encoding GCREC and introducing these vectors by mechanical means into GCREC-deficient cells. Mechanical transfer technologies for

35 use with cells in vivo or ex vitro include (i) direct DNA microinjection into individual cells, (ii)

ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) *Annu. Rev. Biochem.* 62:191-217; Ivics, Z. (1997) *Cell* 91:501-510; Boulay, J-L. and H. Récipon (1998) *Curr. Opin. Biotechnol.* 9:445-450).

- 5 Expression vectors that may be effective for the expression of GCREC include, but are not limited to, the PCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX, PCR2-TOPOTA vectors (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). GCREC may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV),
- 10 Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or β -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551; Gossen, M. et al. (1995) *Science* 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) *Curr. Opin. Biotechnol.* 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND;
- 15 Invitrogen); the FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and Blau, H.M. supra), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding GCREC from a normal individual.

- Commercially available liposome transformation kits (e.g., the PERFECT LIPID TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver
- 20 polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and A.J. Eb (1973) *Virology* 52:456-467), or by electroporation (Neumann, E. et al. (1982) *EMBO J.* 1:841-845). The introduction of DNA to primary cells requires modification of these standardized mammalian transfection protocols.

- 25 In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to GCREC expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding GCREC under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences
- 30 required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) *Proc. Natl. Acad. Sci. USA* 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al.
- 35 (1987) *J. Virol.* 61:1647-1650; Bender, M.A. et al. (1987) *J. Virol.* 61:1639-1646; Adam, M.A. and

A.D. Miller (1988) *J. Virol.* 62:3802-3806; Dull, T. et al. (1998) *J. Virol.* 72:8463-8471; Zufferey, R. et al. (1998) *J. Virol.* 72:9873-9880). U.S. Patent Number 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant") discloses a method for obtaining retrovirus packaging cell lines and is hereby incorporated by
5 reference. Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4⁺ T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) *J. Virol.* 71:7020-7029; Bauer, G. et al. (1997) *Blood* 89:2259-2267; Bonyhadi, M.L. (1997) *J. Virol.* 71:4707-4716; Ranga, U. et al. (1998) *Proc. Natl. Acad. Sci. USA* 95:1201-1206; Su, L. (1997) *Blood* 89:2283-
10 2290).

In the alternative, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding GCREC to cells which have one or more genetic abnormalities with respect to the expression of GCREC. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to
15 be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) *Transplantation* 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent Number 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) *Annu. Rev. Nutr.* 19:511-544 and Verma, I.M. and N. Somia (1997) *Nature* 18:389:239-242, both
20 incorporated by reference herein.

In another alternative, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding GCREC to target cells which have one or more genetic abnormalities with respect to the expression of GCREC. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing GCREC to cells of the central nervous system, for which HSV has
25 a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) *Exp. Eye Res.* 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent Number 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is
30 hereby incorporated by reference. U.S. Patent Number 5,804,413 teaches the use of recombinant HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999) *J. Virol.* 73:519-532 and Xu, H. et al.
35 (1994) *Dev. Biol.* 163:152-161, hereby incorporated by reference. The manipulation of cloned

herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary skill in the art.

5 In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to deliver polynucleotides encoding GCREC to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) *Curr. Opin. Biotechnol.* 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid
10 proteins. This subgenomic RNA replicates to higher levels than the full length genomic RNA, resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g., protease and polymerase). Similarly, inserting the coding sequence for GCREC into the alphavirus genome in place of the capsid-coding region results in the production of a large number of GCREC-coding RNAs and the synthesis of high levels of GCREC in vector transduced cells. While
15 alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application (Dryga, S.A. et al. (1997) *Virology* 228:74-83). The wide host range of alphaviruses will allow the introduction of GCREC into a variety of cell types. The specific transduction of a subset of
20 cells in a population may require the sorting of cells prior to transduction. The methods of manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions
25 -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E.
30 and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme
35 molecule to complementary target RNA, followed by endonucleolytic cleavage. For example,

engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding GCREC.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding GCREC. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding GCREC. Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased GCREC expression or activity, a compound which specifically inhibits expression of the polynucleotide encoding GCREC may be therapeutically useful, and in the treatment of disorders

associated with decreased GCREC expression or activity, a compound which specifically promotes expression of the polynucleotide encoding GCREC may be therapeutically useful.

At least one, and up to a plurality, of test compounds may be screened for effectiveness in altering expression of a specific polynucleotide. A test compound may be obtained by any method commonly known in the art, including chemical modification of a compound known to be effective in
5 altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound based on chemical and/or structural properties of the target polynucleotide; and selection from a library of chemical compounds created combinatorially or randomly. A sample comprising a
10 polynucleotide encoding GCREC is exposed to at least one test compound thus obtained. The sample may comprise, for example, an intact or permeabilized cell, or an in vitro cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding GCREC are assayed by any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence complementary to the sequence
15 of the polynucleotide encoding GCREC. The amount of hybridization may be quantified, thus forming the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide exposed to a test compound indicates that the test compound is effective in altering the expression of the polynucleotide. A screen for a compound effective in altering expression of a specific
20 polynucleotide can be carried out, for example, using a Schizosaccharomyces pombe gene expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide
25 nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Bruce, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruce, T.W. et al. (2000) U.S. Patent No. 6,022,691).

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells
30 taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of

such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a composition which generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient. Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such compositions may consist of GCREC, antibodies to GCREC, and mimetics, agonists, antagonists, or inhibitors of GCREC.

The compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

Compositions for pulmonary administration may be prepared in liquid or dry powder form. These compositions are generally aerosolized immediately prior to inhalation by the patient. In the case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle injection, and obviates the need for potentially toxic penetration enhancers.

Compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

Specialized forms of compositions may be prepared for direct intracellular delivery of macromolecules comprising GCREC or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, GCREC or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et al. (1999) Science 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and

routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example GCREC or fragments thereof, antibodies of GCREC, and agonists, antagonists or inhibitors of GCREC, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about $0.1 \mu\text{g}$ to $100,000 \mu\text{g}$, up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind GCREC may be used for the diagnosis of disorders characterized by expression of GCREC, or in assays to monitor patients being treated with GCREC or agonists, antagonists, or inhibitors of GCREC. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for GCREC include methods which utilize the antibody and a label to detect GCREC in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in

the art and may be used.

A variety of protocols for measuring GCREC, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of GCREC expression. Normal or standard values for GCREC expression are established by combining body fluids or cell
5 extracts taken from normal mammalian subjects, for example, human subjects, with antibodies to GCREC under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of GCREC expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for
10 diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding GCREC may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of GCREC may be correlated
15 with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of GCREC, and to monitor regulation of GCREC levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding GCREC or closely related molecules may be used to identify nucleic acid sequences which encode GCREC. The specificity of the probe, whether it is
20 made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding GCREC, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50%
25 sequence identity to any of the GCREC encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:20-38 or from genomic sequences including promoters, enhancers, and introns of the GCREC gene.

Means for producing specific hybridization probes for DNAs encoding GCREC include the cloning of polynucleotide sequences encoding GCREC or GCREC derivatives into vectors for the
30 production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ³²P or ³⁵S, or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

35 Polynucleotide sequences encoding GCREC may be used for the diagnosis of disorders

associated with expression of GCREC. Examples of such disorders include, but are not limited to, a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including

5 adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, colon, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms,

10 Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous

15 system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial

20 nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic

25 neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; a cardiovascular disorder such as arteriovenous fistula, atherosclerosis, hypertension, vasculitis, Raynaud's disease, aneurysms, arterial dissections, varicose veins, thrombophlebitis and phlebothrombosis, vascular tumors, complications of thrombolysis, balloon angioplasty, vascular replacement, and coronary artery bypass graft surgery, congestive heart failure,

30 ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve, mitral annular calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis, nonbacterial thrombotic endocarditis, endocarditis of systemic lupus erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart

disease, congenital heart disease, and complications of cardiac transplantation; a gastrointestinal disorder such as dysphagia, peptic esophagitis, esophageal spasm, esophageal stricture, esophageal carcinoma, dyspepsia, indigestion, gastritis, gastric carcinoma, anorexia, nausea, emesis, gastroparesis, antral or pyloric edema, abdominal angina, pyrosis, gastroenteritis, intestinal

5 obstruction, infections of the intestinal tract, peptic ulcer, cholelithiasis, cholecystitis, cholestasis, pancreatitis, pancreatic carcinoma, biliary tract disease, hepatitis, hyperbilirubinemia, cirrhosis, passive congestion of the liver, hepatoma, infectious colitis, ulcerative colitis, ulcerative proctitis, Crohn's disease, Whipple's disease, Mallory-Weiss syndrome, colonic carcinoma, colonic

10 obstruction, irritable bowel syndrome, short bowel syndrome, diarrhea, constipation, gastrointestinal hemorrhage, acquired immunodeficiency syndrome (AIDS) enteropathy, jaundice, hepatic encephalopathy, hepatorenal syndrome, hepatic steatosis, hemochromatosis, Wilson's disease, alpha₁-antitrypsin deficiency, Reye's syndrome, primary sclerosing cholangitis, liver infarction, portal vein obstruction and thrombosis, centrilobular necrosis, peliosis hepatis, hepatic vein thrombosis, veno-

15 occlusive disease, preeclampsia, eclampsia, acute fatty liver of pregnancy, intrahepatic cholestasis of pregnancy, and hepatic tumors including nodular hyperplasias, adenomas, and carcinomas; an autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis,

20 cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis,

25 polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a metabolic disorder such as diabetes, obesity, and osteoporosis; and an infection by a viral

30 agent classified as adenovirus, arenavirus, bunyavirus, calicivirus, coronavirus, filovirus, hepadnavirus, herpesvirus, flavivirus, orthomyxovirus, parvovirus, papovavirus, paramyxovirus, picornavirus, poxvirus, reovirus, retrovirus, rhabdovirus, and togavirus. The polynucleotide sequences encoding GCREC may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like

assays; and in microarrays utilizing fluids or tissues from patients to detect altered GCREC expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding GCREC may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding GCREC may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding GCREC in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of GCREC, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding GCREC, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding GCREC may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide

encoding GCREC, or a fragment of a polynucleotide complementary to the polynucleotide encoding GCREC, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

5 In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences encoding GCREC may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP,
10 oligonucleotide primers derived from the polynucleotide sequences encoding GCREC are used to amplify DNA using the polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable using gel electrophoresis in non-denaturing gels. In fSSCP, the
15 oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-throughput equipment such as DNA sequencing machines. Additionally, sequence database analysis methods, termed *in silico* SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation
20 of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

Methods which may also be used to quantify the expression of GCREC include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from
25 standard curves. (See, e.g., Melby, P.C. et al. (1993) *J. Immunol. Methods* 159:235-244; Duplaa, C. et al. (1993) *Anal. Biochem.* 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

30 In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large numbers of genes simultaneously as described below. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene
35 function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor

progression/regression of disease as a function of gene expression, and to develop and monitor the activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

In another embodiment, GCREC, fragments of GCREC, or antibodies specific for GCREC may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by quantifying the number of expressed genes and their relative abundance under given conditions and at a given time. (See Seilhamer et al., "Comparative Gene Transcript Analysis," U.S. Patent Number 5,840,484, expressly incorporated by reference herein.) Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the hybridization takes place in high-throughput format, wherein the polynucleotides of the present invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines, biopsies, or other biological samples. The transcript image may thus reflect gene expression in vivo, as in the case of a tissue or biopsy sample, or in vitro, as in the case of a cell line.

Transcript images which profile the expression of the polynucleotides of the present invention may also be used in conjunction with in vitro model systems and preclinical evaluation of pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental compounds. All compounds induce characteristic gene expression patterns, frequently termed molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and toxicity (Nuwaysir, E.F. et al. (1999) Mol. Carcinog. 24:153-159; Steiner, S. and N.L. Anderson (2000) Toxicol. Lett. 112-113:467-471, expressly incorporated by reference herein). If a test compound has a signature similar to that of a compound with known toxicity, it is likely to share those toxic properties. These fingerprints or signatures are most useful and refined when they contain expression information from a large number of genes and gene families. Ideally, a genome-wide measurement of expression provides the highest quality signature. Even genes whose expression is

not altered by any tested compounds are important as well, as the levels of expression of these genes are used to normalize the rest of the expression data. The normalization procedure is useful for comparison of expression data after treatment with different compounds. While the assignment of gene function to elements of a toxicant signature aids in interpretation of toxicity mechanisms, knowledge of gene function is not necessary for the statistical matching of signatures which leads to prediction of toxicity. (See, for example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released February 29, 2000, available at <http://www.niehs.nih.gov/oc/news/toxchip.htm>.) Therefore, it is important and desirable in toxicological screening using toxicant signatures to include all expressed gene sequences.

10 In one embodiment, the toxicity of a test compound is assessed by treating a biological sample containing nucleic acids with the test compound. Nucleic acids that are expressed in the treated biological sample are hybridized with one or more probes specific to the polynucleotides of the present invention, so that transcript levels corresponding to the polynucleotides of the present invention may be quantified. The transcript levels in the treated biological sample are compared with levels in an untreated biological sample. Differences in the transcript levels between the two samples are indicative of a toxic response caused by the test compound in the treated sample.

Another particular embodiment relates to the use of the polypeptide sequences of the present invention to analyze the proteome of a tissue or cell type. The term proteome refers to the global pattern of protein expression in a particular tissue or cell type. Each protein component of a proteome can be subjected individually to further analysis. Proteome expression patterns, or profiles, are analyzed by quantifying the number of expressed proteins and their relative abundance under given conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and analyzing the polypeptides of a particular tissue or cell type. In one embodiment, the separation is achieved using two-dimensional gel electrophoresis, in which proteins from a sample are separated by isoelectric focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, *supra*). The proteins are visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent such as Coomassie Blue or silver or fluorescent stains. The optical density of each protein spot is generally proportional to the level of the protein in the sample. The optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard methods employing chemical or enzymatic cleavage followed by mass spectrometry. The identity of the protein in a spot may be determined by comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the

polypeptide sequences of the present invention. In some cases, further sequence data may be obtained for definitive protein identification.

A proteomic profile may also be generated using antibodies specific for GCREC to quantify the levels of GCREC expression. In one embodiment, the antibodies are used as elements on a
5 microarray, and protein expression levels are quantified by exposing the microarray to the sample and detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) Anal. Biochem. 270:103-111; Mendoze, L.G. et al. (1999) Biotechniques 27:778-788). Detection may be performed by a variety of methods known in the art, for example, by reacting the proteins in the sample with a thiol- or amino-reactive fluorescent compound and detecting the amount of fluorescence bound at
10 each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor correlation between transcript and protein abundances for some proteins in some tissues (Anderson, N.L. and J. Seilhamer (1997) Electrophoresis 18:533-537), so proteome toxicant signatures may be
15 useful in the analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

In another embodiment, the toxicity of a test compound is assessed by treating a biological
20 sample containing proteins with the test compound. Proteins that are expressed in the treated biological sample are separated so that the amount of each protein can be quantified. The amount of each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample. Individual proteins are identified by sequencing
25 the amino acid residues of the individual proteins and comparing these partial sequences to the polypeptides of the present invention.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins from the biological sample are incubated with antibodies specific to the polypeptides of the present invention. The amount of
30 protein recognized by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g.,
35 Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci.

USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are well known and thoroughly described in DNA Microarrays: A Practical Approach, M. Schena, ed. 5 (1999) Oxford University Press, London, hereby expressly incorporated by reference.

In another embodiment of the invention, nucleic acid sequences encoding GCREC may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among 10 members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. 15 et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism (RFLP). (See, for example, Lander, E.S. and D. Botstein (1986) Proc. Natl. Acad. Sci. USA 83:7353- 20 7357.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding GCREC on a 25 physical map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, 30 may reveal associated markers even if the exact chromosomal locus is not known. This information is valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further 35 investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of

the instant invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, GCREC, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between GCREC and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with GCREC, or fragments thereof, and washed. Bound GCREC is then detected by methods well known in the art. Purified GCREC can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding GCREC specifically compete with a test compound for binding GCREC. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with GCREC.

In additional embodiments, the nucleotide sequences which encode GCREC may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. No. 60/221,478, U.S. Ser. No. 60/223,268, U.S. Ser. No. 60/231,121, U.S. Ser. No. 60/232,691, U.S. Ser. No. 60/235,146, U.S. Ser. No. 60/227,054, and U.S. Ser. No. 60/232,243, are hereby expressly incorporated by reference.

EXAMPLES

I. Construction of cDNA Libraries

Incyte cDNAs were derived from cDNA libraries described in the LIFESEQ GOLD database

(Incyte Genomics, Palo Alto CA) and shown in Table 4, column 5. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or
5 extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A)+ RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN,
10 Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP
15 vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-
20 1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), PCDNA2.1 plasmid (Invitrogen, Carlsbad CA), PBK-CMV plasmid (Stratagene), or pINCY (Incyte Genomics, Palo Alto
25 CA), or derivatives thereof. Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

Plasmids obtained as described in Example I were recovered from host cells by in vivo
30 excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1
35 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows. Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Applied Biosystems) thermal cyclers or the PTC-200 thermal cyclers (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (Applied Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, *supra*, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VIII.

The polynucleotide sequences derived from Incyte cDNAs were validated by removing vector, linker, and poly(A) sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The Incyte cDNA sequences or translations thereof were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and hidden Markov model (HMM)-based protein family databases such as PFAM. (HMM is a probabilistic approach which analyzes consensus primary structures of gene families. See, for example, Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.) The queries were performed using programs based on BLAST, FASTA, BLIMPS, and HMMER. The Incyte cDNA sequences were assembled to produce full length polynucleotide sequences. Alternatively, GenBank cDNAs, GenBank ESTs, stitched sequences, stretched sequences, or Genscan-predicted coding sequences (see Examples IV and V) were used to extend Incyte cDNA assemblages to full length. Assembly was performed using programs based on Phred, Phrap, and Consed, and cDNA assemblages were screened for open reading frames using programs based on

GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length polypeptide sequences. Alternatively, a polypeptide of the invention may begin at any of the methionine residues of the full length translated polypeptide. Full length polypeptide sequences were subsequently analyzed by querying against databases such as the

5 GenBank protein databases (genpept), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and hidden Markov model (HMM)-based protein family databases such as PFAM. Full length polynucleotide sequences are also analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments are generated using default parameters specified by the

10 CLUSTAL algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

Table 7 summarizes the tools, programs, and algorithms used for the analysis and assembly of Incyte cDNA and full length sequences and provides applicable descriptions, references, and threshold parameters. The first column of Table 7 shows the tools, programs, and algorithms used,

15 the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score or the lower the probability value, the greater the identity between two sequences).

20 The programs described above for the assembly and analysis of full length polynucleotide and polypeptide sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:20-38. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies are described in Table 4, column 4.

IV. Identification and Editing of Coding Sequences from Genomic DNA

25 Putative G-protein coupled receptors were initially identified by running the Genscan gene identification program against public genomic sequence databases (e.g., gbpri and gbhtg). Genscan is a general-purpose gene identification program which analyzes genomic DNA sequences from a variety of organisms (See Burge, C. and S. Karlin (1997) J. Mol. Biol. 268:78-94, and Burge, C. and S. Karlin (1998) Curr. Opin. Struct. Biol. 8:346-354). The program concatenates predicted exons to

30 form an assembled cDNA sequence extending from a methionine to a stop codon. The output of Genscan is a FASTA database of polynucleotide and polypeptide sequences. The maximum range of sequence for Genscan to analyze at once was set to 30 kb. To determine which of these Genscan predicted cDNA sequences encode G-protein coupled receptors, the encoded polypeptides were analyzed by querying against PFAM models for G-protein coupled receptors. Potential G-protein

35 coupled receptors were also identified by homology to Incyte cDNA sequences that had been

annotated as G-protein coupled receptors. These selected Genscan-predicted sequences were then compared by BLAST analysis to the genpept and gbpr public databases. Where necessary, the Genscan-predicted sequences were then edited by comparison to the top BLAST hit from genpept to correct errors in the sequence predicted by Genscan, such as extra or omitted exons. BLAST analysis was also used to find any Incyte cDNA or public cDNA coverage of the Genscan-predicted sequences, thus providing evidence for transcription. When Incyte cDNA coverage was available, this information was used to correct or confirm the Genscan predicted sequence. Full length polynucleotide sequences were obtained by assembling Genscan-predicted coding sequences with Incyte cDNA sequences and/or public cDNA sequences using the assembly process described in Example III. Alternatively, full length polynucleotide sequences were derived entirely from edited or unedited Genscan-predicted coding sequences.

V. Assembly of Genomic Sequence Data with cDNA Sequence Data

"Stitched" Sequences

Partial cDNA sequences were extended with exons predicted by the Genscan gene identification program described in Example IV. Partial cDNAs assembled as described in Example III were mapped to genomic DNA and parsed into clusters containing related cDNAs and Genscan exon predictions from one or more genomic sequences. Each cluster was analyzed using an algorithm based on graph theory and dynamic programming to integrate cDNA and genomic information, generating possible splice variants that were subsequently confirmed, edited, or extended to create a full length sequence. Sequence intervals in which the entire length of the interval was present on more than one sequence in the cluster were identified, and intervals thus identified were considered to be equivalent by transitivity. For example, if an interval was present on a cDNA and two genomic sequences, then all three intervals were considered to be equivalent. This process allows unrelated but consecutive genomic sequences to be brought together, bridged by cDNA sequence. Intervals thus identified were then "stitched" together by the stitching algorithm in the order that they appear along their parent sequences to generate the longest possible sequence, as well as sequence variants. Linkages between intervals which proceed along one type of parent sequence (cDNA to cDNA or genomic sequence to genomic sequence) were given preference over linkages which change parent type (cDNA to genomic sequence). The resultant stitched sequences were translated and compared by BLAST analysis to the genpept and gbpr public databases. Incorrect exons predicted by Genscan were corrected by comparison to the top BLAST hit from genpept. Sequences were further extended with additional cDNA sequences, or by inspection of genomic DNA, when necessary.

"Stretched" Sequences

Partial DNA sequences were extended to full length with an algorithm based on BLAST analysis. First, partial cDNAs assembled as described in Example III were queried against public

databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases using the BLAST program. The nearest GenBank protein homolog was then compared by BLAST analysis to either Incyte cDNA sequences or GenScan exon predicted sequences described in Example IV. A chimeric protein was generated by using the resultant high-scoring segment pairs (HSPs) to map the translated sequences onto the GenBank protein homolog. Insertions or deletions may occur in the chimeric protein with respect to the original GenBank protein homolog. The GenBank protein homolog, the chimeric protein, or both were used as probes to search for homologous genomic sequences from the public human genome databases. Partial DNA sequences were therefore "stretched" or extended by the addition of homologous genomic sequences. The resultant stretched sequences were examined to determine whether it contained a complete gene.

VI. Chromosomal Mapping of GCREC Encoding Polynucleotides

The sequences which were used to assemble SEQ ID NO:20-38 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched SEQ ID NO:20-38 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 7). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO., to that map location.

Map locations are represented by ranges, or intervals, of human chromosomes. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Généthon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Human genome maps and other resources available to the public, such as the NCBI "GeneMap'99" World Wide Web site (<http://www.ncbi.nlm.nih.gov/genemap/>), can be employed to determine if previously identified disease genes map within or in proximity to the intervals indicated above.

VII. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, *supra*, ch. 7; Ausubel (1995) *supra*, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum} \{ \text{length}(\text{Seq. 1}), \text{length}(\text{Seq. 2}) \}}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

Alternatively, polynucleotide sequences encoding GCREC are analyzed with respect to the tissue sources from which they were derived. For example, some full length sequences are assembled, at least in part, with overlapping Incyte cDNA sequences (see Example III). Each cDNA sequence is derived from a cDNA library constructed from a human tissue. Each human tissue is classified into one of the following organ/tissue categories: cardiovascular system; connective tissue; digestive system; embryonic structures; endocrine system; exocrine glands; genitalia, female; genitalia, male; germ cells; hemic and immune system; liver; musculoskeletal system; nervous system; pancreas; respiratory system; sense organs; skin; stomatognathic system; unclassified/mixed; or urinary tract. The number of libraries in each category is counted and divided by the total number of libraries across all categories. Similarly, each human tissue is classified into one of the following disease/condition categories: cancer, cell line, developmental, inflammation, neurological, trauma, cardiovascular, pooled, and other, and the number of libraries in each category is counted and divided by the total number of libraries across all categories. The resulting percentages reflect the tissue- and

disease-specific expression of cDNA encoding GCREC. cDNA sequences and cDNA library/tissue information are found in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA).

VIII. Extension of GCREC Encoding Polynucleotides

Full length polynucleotide sequences were also produced by extension of an appropriate
5 fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer was synthesized to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target
10 sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR
15 was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and 2-mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the
20 alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100 μ l PICOGREEN
25 quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 μ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by
30 electrophoresis on a 1 % agarose gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviII cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For
35 shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%)

agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on
5 antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min;
10 Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM
15 BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems).

In like manner, full length polynucleotide sequences are verified using the above procedure or are used to obtain 5' regulatory sequences using the above procedure along with oligonucleotides designed for such extension, and an appropriate genomic library.

IX. Labeling and Use of Individual Hybridization Probes

20 Hybridization probes derived from SEQ ID NO:20-38 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 µCi of
25 [γ-³²P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10⁷ counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases:
30 Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate.
35 Hybridization patterns are visualized using autoradiography or an alternative imaging means and

compared.

X. Microarrays

The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschweiler, supra), mechanical microspotting technologies, and derivatives thereof. The substrate in each of the
5 aforementioned technologies should be uniform and solid with a non-porous surface (Schena (1999), supra). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers. Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding
10 procedures. A typical array may be produced using available methods and machines well known to those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g., Schena, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645; Marshall, A. and J. Hodgson (1998) Nat. Biotechnol. 16:27-31.)

Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may
15 comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection. After hybridization, nonhybridized nucleotides from the biological sample are removed, and a
20 fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

25 Tissue or Cell Sample Preparation

Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and poly(A)⁺ RNA is purified using the oligo-(dT) cellulose method. Each poly(A)⁺ RNA sample is reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/ μ l oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/ μ l RNase inhibitor, 500 μ M dATP, 500 μ M dGTP, 500 μ M dTTP, 40
30 μ M dCTP, 40 μ M dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)⁺ RNA with GEMBRIGHT kits (Incyte). Specific control poly(A)⁺ RNAs are synthesized by in vitro transcription from non-coding yeast genomic DNA. After incubation at 37°C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and
35 incubated for 20 minutes at 85°C to stop the reaction and degrade the RNA. Samples are purified

using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc. (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and
5 resuspended in 14 μ l 5X SSC/0.2% SDS.

Microarray Preparation

Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are
10 amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5 μ g. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia Biotech).

Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water
15 washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a 110°C oven.

Array elements are applied to the coated glass substrate using a procedure described in US
20 Patent No. 5,807,522, incorporated herein by reference. 1 μ l of the array element DNA, at an average concentration of 100 ng/ μ l, is loaded into the open capillary printing element by a high-speed robotic apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water.
25 Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60°C followed by washes in 0.2% SDS and distilled water as before.

Hybridization

Hybridization reactions contain 9 μ l of sample mixture consisting of 0.2 μ g each of Cy3 and
30 Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65°C for 5 minutes and is aliquoted onto the microarray surface and covered with an 1.8 cm² coverslip. The arrays are transferred to a waterproof chamber having a cavity just slightly larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140 μ l of 5X SSC in a corner of the chamber. The chamber containing the arrays is

incubated for about 6.5 hours at 60°C. The arrays are washed for 10 min at 45°C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45°C in a second wash buffer (0.1X SSC), and dried.

Detection

5 Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477, Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate
15 filters positioned between the array and the photomultiplier tubes are used to filter the signals. The emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

The sensitivity of the scans is typically calibrated using the signal intensity generated by a
20 cDNA control species added to the sample mixture at a known concentration. A specific location on the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different fluorophore, are hybridized to a single array for the purpose of identifying genes that are
25 differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC computer. The digitized data are displayed as an image where the signal intensity is mapped using a
30 linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

A grid is superimposed over the fluorescence signal image such that the signal from each

spot is centered in each element of the grid. The fluorescence signal within each element is then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

XI. Complementary Polynucleotides

- 5 Sequences complementary to the GCREC-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring GCREC. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of GCREC.
- 10 To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the GCREC-encoding transcript.

XII. Expression of GCREC

- 15 Expression and purification of GCREC is achieved using bacterial or virus-based expression systems. For expression of GCREC in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory
- 20 element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express GCREC upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of GCREC in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is
- 25 replaced with cDNA encoding GCREC by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K.
- 30 et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

- In most expression systems, GCREC is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-
- 35 kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on

immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from GCREC at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified GCREC obtained by these methods can be used directly in the assays shown in Examples XVI, XVII, and XVIII, where applicable.

XIII. Functional Assays

GCREC function is assessed by expressing the sequences encoding GCREC at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include PCMV SPORT (Life Technologies) and PCR3.1 (Invitrogen, Carlsbad CA), both of which contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of GCREC on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding GCREC and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake

Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding GCREC and other genes of interest can be analyzed by northern analysis or microarray techniques.

XIV. Production of GCREC Specific Antibodies

5 GCREC substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the GCREC amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is
10 synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, *supra*, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Applied Biosystems) using Fmoc chemistry and coupled to KLH (Sigma-
15 Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, *supra*.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-GCREC activity by, for example, binding the peptide or GCREC to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat
20 anti-rabbit IgG.

XV. Purification of Naturally Occurring GCREC Using Specific Antibodies

Naturally occurring or recombinant GCREC is substantially purified by immunoaffinity chromatography using antibodies specific for GCREC. An immunoaffinity column is constructed by covalently coupling anti-GCREC antibody to an activated chromatographic resin, such as
25 CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing GCREC are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of GCREC (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt
30 antibody/GCREC binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and GCREC is collected.

XVI. Identification of Molecules Which Interact with GCREC

GCREC, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules
35 previously arrayed in the wells of a multi-well plate are incubated with the labeled GCREC, washed,

and any wells with labeled GCREC complex are assayed. Data obtained using different concentrations of GCREC are used to calculate values for the number, affinity, and association of GCREC with the candidate molecules.

Alternatively, molecules interacting with GCREC are analyzed using the yeast two-hybrid
5 system as described in Fields, S. and O. Song (1989) Nature 340:245-246, or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

GCREC may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT) which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S.
10 Patent No. 6,057,101).

XVII. Demonstration of GCREC Activity

An assay for GCREC activity measures the expression of GCREC on the cell surface. cDNA encoding GCREC is transfected into an appropriate mammalian cell line. Cell surface proteins are labeled with biotin as described (de la Fuente, M.A. et al. (1997) Blood 90:2398-2405).
15 Immunoprecipitations are performed using GCREC-specific antibodies, and immunoprecipitated samples are analyzed using sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and immunoblotting techniques. The ratio of labeled immunoprecipitant to unlabeled immunoprecipitant is proportional to the amount of GCREC expressed on the cell surface.

In the alternative, an assay for GCREC activity is based on a prototypical assay for
20 ligand/receptor-mediated modulation of cell proliferation. This assay measures the rate of DNA synthesis in Swiss mouse 3T3 cells. A plasmid containing polynucleotides encoding GCREC is added to quiescent 3T3 cultured cells using transfection methods well known in the art. The transiently transfected cells are then incubated in the presence of [³H]thymidine, a radioactive DNA precursor molecule. Varying amounts of GCREC ligand are then added to the cultured cells.
25 Incorporation of [³H]thymidine into acid-precipitable DNA is measured over an appropriate time interval using a radioisotope counter, and the amount incorporated is directly proportional to the amount of newly synthesized DNA. A linear dose-response curve over at least a hundred-fold GCREC ligand concentration range is indicative of receptor activity. One unit of activity per milliliter is defined as the concentration of GCREC producing a 50% response level, where 100%
30 represents maximal incorporation of [³H]thymidine into acid-precipitable DNA (McKay, I. and I. Leigh, eds. (1993) Growth Factors: A Practical Approach, Oxford University Press, New York NY, p. 73.)

In a further alternative, the assay for GCREC activity is based upon the ability of GPCR family proteins to modulate G protein-activated second messenger signal transduction pathways (e.g.,
35 cAMP; Gaudin, P. et al. (1998) J. Biol. Chem. 273:4990-4996). A plasmid encoding full length

GCREC is transfected into a mammalian cell line (e.g., Chinese hamster ovary (CHO) or human embryonic kidney (HEK-293) cell lines) using methods well-known in the art. Transfected cells are grown in 12-well trays in culture medium for 48 hours, then the culture medium is discarded, and the attached cells are gently washed with PBS. The cells are then incubated in culture medium with or without ligand for 30 minutes, then the medium is removed and cells lysed by treatment with 1 M perchloric acid. The cAMP levels in the lysate are measured by radioimmunoassay using methods well-known in the art. Changes in the levels of cAMP in the lysate from cells exposed to ligand compared to those without ligand are proportional to the amount of GCREC present in the transfected cells.

To measure changes in inositol phosphate levels, the cells are grown in 24-well plates containing 1×10^5 cells/well and incubated with inositol-free media and [^3H]myoinositol, 2 μCi /well, for 48 hr. The culture medium is removed, and the cells washed with buffer containing 10 mM LiCl followed by addition of ligand. The reaction is stopped by addition of perchloric acid. Inositol phosphates are extracted and separated on Dowex AG1-X8 (Bio-Rad) anion exchange resin, and the total labeled inositol phosphates counted by liquid scintillation. Changes in the levels of labeled inositol phosphate from cells exposed to ligand compared to those without ligand are proportional to the amount of GCREC present in the transfected cells.

XVIII. Identification of GCREC Ligands

GCREC is expressed in a eukaryotic cell line such as CHO (Chinese Hamster Ovary) or HEK (Human Embryonic Kidney) 293 which have a good history of GPCR expression and which contain a wide range of G-proteins allowing for functional coupling of the expressed GCREC to downstream effectors. The transformed cells are assayed for activation of the expressed receptors in the presence of candidate ligands. Activity is measured by changes in intracellular second messengers, such as cyclic AMP or Ca^{2+} . These may be measured directly using standard methods well known in the art, or by the use of reporter gene assays in which a luminescent protein (e.g. firefly luciferase or green fluorescent protein) is under the transcriptional control of a promoter responsive to the stimulation of protein kinase C by the activated receptor (Milligan, G. et al. (1996) Trends Pharmacol. Sci. 17:235-237). Assay technologies are available for both of these second messenger systems to allow high throughput readout in multi-well plate format, such as the adenylyl cyclase activation FlashPlate Assay (NEN Life Sciences Products), or fluorescent Ca^{2+} indicators such as Fluo-4 AM (Molecular Probes) in combination with the FLIPR fluorimetric plate reading system (Molecular Devices). In cases where the physiologically relevant second messenger pathway is not known, GCREC may be coexpressed with the G-proteins $G_{\alpha 15/16}$ which have been demonstrated to couple to a wide range of G-proteins (Offermanns, S. and M.L. Simon (1995) J. Biol. Chem. 270:15175-15180), in order to funnel the signal transduction of the GCREC through a pathway involving phospholipase C and Ca^{2+}

mobilization. Alternatively, GCREC may be expressed in engineered yeast systems which lack endogenous GPCRs, thus providing the advantage of a null background for GCREC activation screening. These yeast systems substitute a human GPCR and Ga protein for the corresponding components of the endogenous yeast pheromone receptor pathway. Downstream signaling pathways
5 are also modified so that the normal yeast response to the signal is converted to positive growth on selective media or to reporter gene expression (Broach, J.R. and J. Thorner (1996) Nature 384 (supp.):14-16). The receptors are screened against putative ligands including known GPCR ligands and other naturally occurring bioactive molecules. Biological extracts from tissues, biological fluids and cell supernatants are also screened.

10

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific
15 embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Incyte Project ID	Polypeptide SEQ ID NO:	Incyte Polypeptide ID	Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID
7474806	1	7474806CD1	20	7474806CB1
7474840	2	7474840CD1	21	7474840CB1
7475092	3	7475092CD1	22	7475092CB1
7341260	4	7341260CD1	23	7341260CB1
7473911	5	7473911CD1	24	7473911CB1
7474767	6	7474767CD1	25	7474767CB1
7475815	7	7475815CD1	26	7475815CB1
60263275	8	60263275CD1	27	60263275CB1
60203310	9	60203310CD1	28	60203310CB1
7477349	10	7477349CD1	29	7477349CB1
55002225	11	55002225CD1	30	55002225CB1
7475686	12	7475686CD1	31	7475686CB1
7482007	13	7482007CD1	32	7482007CB1
6769042	14	6769042CD1	33	6769042CB1
7476053	15	7476053CD1	34	7476053CB1
7480410	16	7480410CD1	35	7480410CB1
55036418	17	55036418CD1	36	55036418CB1
7481701	18	7481701CD1	37	7481701CB1
7481774	19	7481774CD1	38	7481774CB1

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank NO:	GenBank ID	Probability score	GenBank Homolog
1	7474806CD1	g2707256		4.00E-62	[fl][Meleagris gallopavo] G protein coupled P2Y nucleotide receptor
2	7474840CD1	g11967419		4.00E-65	[fl][Mus musculus] vomeronasal (pheromone) receptor VIRC3
3	7475092CD1	g2992628		2.00E-86	[fl][Homo sapiens] putative seven pass transmembrane protein
4	7341260CD1	g3805932		1.00E-10	[fl][Homo sapiens] putative G-Protein coupled receptor, EDG6
5	7473911CD1	g1055254		2.00E-18	Graler, M.H. et al. (1998) Genomics 53:164-169
6	7474767CD1	g179985		2.00E-14	[fl][Rattus norvegicus] pheromone receptor VN6
7	7475815CD1	g1055254		1.00E-56	Dulac, C. and Axel, R. (1995) Cell 83: 195-206
8	60263275CD1	g13183149		0	[fl][Homo sapiens] C-C chemokine receptor type 1
9	60203310CD1	g3882981		0	Neote, K. et al. (1993) Cell 72:415-425
10	7477349CD1	g6979162		3.00E-11	[fl][Rattus norvegicus] pheromone receptor VN6
11	55002225CD1	g14164383		0	[fl][Homo sapiens] (AF239764) EGF-like module-containing mucin-like receptor EMR3
12	7475686CD1	g7248884		1.00E-180	[fl][Rattus norvegicus] calcium-independent alpha-latrotoxin receptor
13	7482007CD1	g5525078		1.00E-104	[fl][Rattus norvegicus] macrophage inflammatory protein-1 alpha receptor
14	6769042CD1	g4164061		1.70E-59	[fl][Homo sapiens] (AB060151) G protein-coupled receptor
					[fl][Mus musculus] G-protein coupled receptor GPR73
					Parker, R. et al. (2000) Biochim. Biophys. Acta 1491:369-375
					[fl][Rattus norvegicus] seven transmembrane receptor
					Abe, J. et al. (1999) J. Biol. Chem. 274:19957-19964
					[Bos taurus] latrophilin 3 splice variant abbg
					FEBS Lett. (1999) 443:348-352

Table 2 (cont.)

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability score	GenBank Homolog
15	7476053CD1	g310075	1.00E-165	[fl][Rattus norvegicus] serotonin receptor Erlander, M.G. et al. (1993) Proc. Natl. Acad. Sci. U.S.A. 90:3452-3456
16	7480410CD1	g3983382	1.90E-88	[Mus musculus] olfactory receptor E3
17	55036418CD1	g3983398	5.00E-99	[Mus musculus] olfactory receptor G3 Krautwurst, D. et al. (1998) Cell 95:917-926
18	7481701CD1	g12007416	2.00E-67	[fl][Mus musculus] m51 olfactory receptor
19	7481774CD1	g5901478	3.10E-106	[Marmota marmota] olfactory receptor

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
1	7474806CD1	339	S111 S15 S189 S260 S336 S46 S84 T163 T204 T255 T36 T80	N35 N69	Transmembrane domains: Y89-T109, F130-F149 7 transmembrane receptor (rhodopsin family): L68-Y324 G-protein coupled receptor BL00237: W116-P155, F224-Y235, T255-T281, S316-I332 Rhodopsin-like GPCR superfamily PR00237: Y53-W77, A86-I107, F130-V152, H166-A187, G216-V239, S260-F284, Y306-I332 G-protein coupled receptor: DM00013 P41231 27-322: F45-S327 P51582 29-322: F45-S327 I55450 20-317: F45-R328 P48042 45-340: Y53-S327 G-protein coupled receptor: PD000009: Y89-F188 Pheromone receptor: PD009900: I59-M329	HMMER HMMER-PFAM BLIMPS-BLOCKS BLIMPS-PRINTS BLAST-DOMO BLAST-PRODOM BLAST-PRODOM
2	7474840CD1	335	S237 S28 S318 T22 T86	N168 N184	Transmembrane domains: H166-L186, M240-Y256, Y299-L319 Putative seven pass transmembrane protein: PD138976: R3-L360 Transmembrane Domain: L55-L74 7 transmembrane receptor (rhodopsin family): A31-A250 Rhodopsin-like GPCR superfamily signature PR00237B:A50-L71 PR00237C:P92-V114 PR00237D:S123-L144 PR00237F:Q222-L246 PR00237G:L262-T288 Transmembrane Domain: I118-A136 PHEROMONE RECEPTOR PD009900: S87-V167, K42-Y90	HMMER BLAST-PRODOM HMMER HMMER-PFAM BLIMPS-PRINTS HMMER BLAST-PRODOM
3	7475092CD1	428	S257 S286 S295 S342 S356 S365 S412 S79 T156	N180 N207 N284		
4	7341260CD1	330	S123 S185 S323 S5 S78 T219	N154 N4 N76 N93		
5	7473911CD1	676	S105 S213 S215 S250 S330 S353 S492 S569 S591 S606 S646 S647 S79 T217 T296 T301 T394 T620	N184		

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
6	7474767CD1	372	S214 S289 S351 T145 T223 T299	N11 N318 N353	Signal Peptide: M1-T47 Transmembrane Domains: Y32-S52, I74-P100, I184-Y208 7 transmembrane receptor (rhodopsin family): A43-Y285 G-protein coupled receptors proteins BL00237:M94-P133, Y221-M247, N277-R293 Rhodopsin-like GPCR superfamily signature PR00237: V28-S52, S62-V83, E108-V130, R144-W165, I184-V207, T226-Y250, S267-R293 G-protein coupled receptors signature: E105-T155 G-PROTEIN COUPLED RECEPTORS DM00013 P32246 28-316:A22-R293 P51677 28-316:F27-R293 P25930 33-319:F27-L300 P34981 19-336:V28-I206, S222-R293 Transmembrane Domain: I216-A234 PHEROMONE RECEPTOR PD009900: K42-S170, I112-K271	SPSCAN HMMER HMMER_PFAM BLIMPS_BLOCKS BLIMPS_PRINTS PROFILESCAN BLAST_DOMO HMMER BLAST_PRODOM
7	7475815CD1	271	S115 S142 S188 S203 S79 T269			

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
8	60263275CD1	611	S257 S308 S494 S578 S580 S588 S598 S63 S81 T21 T211 T258 T307 T406 T75 Y585	N104 N148 N161 N209 N238 N286 N293 N345 N409 N414	Signal Peptide: M1-Q22 Transmembrane Domains: V321-L339, W424-T451, M464-I488, Y540-L560 7 transmembrane receptor (Secretin family): D312-V564 Latrophilin/CL-1-like GPS domain: K259-Q309 EGF-like domain: C30-C76 G-protein coupled receptor BL00649: C378-L403, G430-R454, W465-S494 Secretin-like GPCR superfamily signature PR00249: V317-K341, I380-L403, K423-L448 W465-K490, A539-L560 Type I EGF signature PR00009: K25-F40, E48-Y59 G-protein coupled receptors family 2 signatures: V531-E574 RECEPTOR G PROTEIN COUPLED PD000752: I315-F572 RECEPTOR G PROTEIN COUPLED EGF LIKE PD005428: N176-M304 HORMONE; EMR1; LEUCOCYTE; ANTIGEN; DM05221 A57172 465-886:S183-K596 I37225 347-738:K259-G592 P48960 347-738:K259-G592 G-PROTEIN COUPLED RECEPTORS FAMILY 2 DM00378 I49149 56-425:V314-R573, G9-V56 Aspartic acid and asparagine hydroxylation site: C44-C55	HMMER HMMER HMMER_PFAM HMMER_PFAM HMMER_PFAM BLIMPS_BLOCKS BLIMPS_PRINTS BLIMPS_PRINTS PROFILES SCAN BLAST_PROD OM BLAST_PROD OM BLAST_DOMO BLAST_DOMO MOTIFS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
9	60203310CD1	1469	S166 S186 S1214 S187 S261 T1253 S325 S351 S1295 S417 S423 T1398 S460 S591 T1453 S630 S761 T1008 S791 S824 S1144 S88 S907 S1154 S978 T113 T1181 T181 T199 S1295 T294 T303 T1346 T443 T559 S1458 T658 T672 T731 T803 T842 Y121	N464 N549 N759 N772 N817 N843 N93 N932 N1098 N1179 N1196 N1251	Signal peptide: M1-S21 Transmembrane domains: V879-F897, V942-L960, F1020-G1040, F1071-W1090 7 transmembrane receptor (secretin family): D874-V1130 Latrophilin/CL-1-like GPCR domain: F814-V866 G-protein coupled receptor BL00649: C940-L965, G987-V1011, W1022-A1051, S1113-L1138, C498-T525, G884-I929 Secretin-like GPCR superfamily PR00249: V879-R903, V942-L965, R980-S1005, W1022-V1047, A1105-L1126 EMRI hormone leucocyte antigen: DM05221 I37225 347-738: L769-S1160 P48960 347-738: L769-S1160 A57172 465-886: N730-T1162 G-protein coupled receptors family 2: DM00378 I49149 56-425: G836-R1131 Latrophilin-related receptor: PD024331: K386-E628 PD041747: L1207-L1469 Myocilin/Olfactomedin response protein: PD006897: Y177-V385 EGF-like G-protein coupled receptor: PD005428: R629-V866 Transmembrane domains: P166-L188, V215-V233 7 transmembrane receptor (rhodopsin family): A251-Y421 G-protein coupled receptor motif: A251-L267 G-protein coupled receptors signature: I243-A293 G-protein coupled receptor BL00237: R231-P270, Y329-N340, P357-V383, H413-R429 Rhodopsin-like GPCR superfamily: I165-A189, S199-A220, E245-L267, R281-W302, K321-I344, S362-Y386, L403-R429 G-protein coupled receptor: DM00013 I49339 28-316: A162-V435 P51675 28-317: A162-V435	HMMER HMMER HMMER-PFAM HMMER-PFAM BLIMPS-BLOCKS BLIMPS-PRINTS BLAST-DOMO BLAST-PRODROM BLAST-PRODROM BLAST-PRODROM HMMER HMMER-PFAM MOTIFS ProfileScan BLIMPS-BLOCKS BLIMPS-PRINTS BLAST-DOMO BLAST-DOMO
10	7477349CD1	469	S122 S199 S25 S312 S447 S85 T110 T195 T207 T282 T315 T39 T427 T431 T465	N139		

Table 3 (cont.)

SEQ ID NO.	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
11	55002225CD1	335	S190 S30 S62 T11 T110 T150 T189 T320 T330 T66	N10 N17	G-PROTEIN COUPLED RECEPTORS DM00013 P32745 38-329:V32-S310 RECEPTOR COUPLED G-PROTEIN TRANSMEMBRANE GLYCOPROTEIN PHOSPHORYLATION LIPOPROTEIN PALMITATE PROTEIN FAMILY PD000009:R61-Y171 G-protein coupled receptor BL00237:S298-Q314, W99-P138, F206-Y217, Q244-I270 Rhodopsin-like GPCR superfamily signature PR00237:I36-I60, P68-L89, D113-L135, K149-V170, T198-L221, L249-V273, Y288-Q314 Transmembrane domain transmem_domain:T34-F57, V194-I218, M252-L272 7 transmembrane receptor (rhodopsin family) 7tm_1:G51-Y306 G-Protein Receptor A119-L135 g-protein coupled receptors signature g_protein_receptor.prf:T110-L157 G-PROTEIN COUPLED RECEPTORS DM00013 P49146 44-341:K299-F590 RECEPTOR COUPLED G-PROTEIN TRANSMEMBRANE GLYCOPROTEIN PHOSPHORYLATION LIPOPROTEIN PALMITATE PROTEIN FAMILY PD000009:K328-F437 G-protein coupled receptor BL00237:W367-P406, F478-Y489, R514-F540 Rhodopsin-like GPCR superfamily signature PR00237:R381-I403, T415-Y436, F470-S493, T519-V543, F561-M587, V301-T325, T334-F355. Neuropeptide Y receptor PR01012:R326-I338, R381-A396 Transmembrane domain transmem_domain: I302-F320 7 transmembrane receptor (rhodopsin family) 7tm_1:G316-N575 G-Protein Receptor V387-I403 G-protein coupled receptors signature g_protein_receptor.prf:N378-I428	BLAST_DOMO BLAST_PRODUM BLIMPS_BLOCKS BLIMPS_PRINTS HMMER HMMER_PPFAM MOTIFS PROFILESSCAN BLAST_DOMO BLAST_PRODUM BLIMPS_BLOCKS BLIMPS_PRINTS BLIMPS_PRINTS HMMER HMMER_PPFAM MOTIFS PROFILESSCAN
12	7475686CD1	630	S230 S269 S601 S607 T223 T519 T581 T586 T620 T81	N332 N584		

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
13	7482007CD1	695	S185 S252 S354 S392 S395 S591 S669 T265 T397 T45 T584 T624 T653	N169 N177 N209 N229 N250 N257 N263 N286 N309 N340 N379 N61 N679 N686	HORMONE; EMR1; LEUCOCYTE; ANTIGEN; DM05221 I37225 347-738:C349-S688 RECEPTOR TRANSMEMBRANE G-PROTEIN COUPLED GLYCOPROTEIN PRECURSOR SIGNAL TYPE POLYPEPTIDE ALTERNATIVE PD000752:N372-R660 G-protein coupled receptors family 2 proteins. BL00649:C473-I498 Secretin-like GPCR superfamily signature PR00249:Y403-W427, A475-I498 Transmembrane domain transmem_domain:I407-T433, M514-T537, L487-F506, L559-V579 7 transmembrane receptor (Secretin family) 7tm_2:D398-I659	BLAST_DOMO BLAST_PRODUM BLIMPS_BLOCKS BLIMPS_PRINTS HMMER HMMER_PPFAM
14	6769042CD1	633	S108 S262 S275 S381 S589 S612 S616 S70 T251 T355 T517	N153 N235 N260 N263 N292 N366 N41 N604 N61 N78	HORMONE; EMR1; LEUCOCYTE; ANTIGEN; DM05221 A57172 465-886:C279-S593 RECEPTOR TRANSMEMBRANE G-PROTEIN COUPLED GLYCOPROTEIN PRECURSOR SIGNAL TYPE POLYPEPTIDE ALTERNATIVE PD000752:H321-K579 G-protein coupled receptors family 2 proteins. BL00649:S555-I580, C391-L416 cAMP-type GPCR signature PR00247:Y361-R383, A395-I421, Y433-S451 Secretin-like GPCR superfamily signature PR00249:S327-S351, V393-L416, H431-S456, W473-S498, T517-L537, Q547-L568 Transmembrane domain transmem_domain:C333-L350, I472-T494 7 transmembrane receptor (Secretin family) 7tm_2:Q322-V572 Latrophilin/CL-1-like GPS domain GPS:V265-L315 G-protein coupled receptors family 2 signatures g_protein_recep_f2_2.prf: A474- S593	BLAST_DOMO BLAST_PRODUM BLIMPS_BLOCKS BLIMPS_PRINTS BLIMPS_PRINTS HMMER HMMER_PPFAM HMMER_PPFAM PROFILES SCAN

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
15	7476053CD1	370	S217 S27 S287 T116 T163		Signal peptide: M298-T317 Signal cleavage: M1-A65 Transmembrane domain: F48-L68, M298-L316 7 transmembrane receptor (rhodopsin family) 7tm_1: W69-Y351 G-protein coupled receptor BL00237: R120-H159, Q291-T317, N343-N359 Rhodopsin-like GPCR superfamily signature PR00237: L54-P78, P87-P108, D134-I156, A170-L191, A214-Y237, A296-I320, K333-N359 5-hydroxytryptamine 5B receptor PR00519: A3-P19, E20-P36, P36-V50, E196-R204, R246-V260, V260-V270 5HYDROXYTRYPTAMINE 5B RECEPTOR 5HT5B SEROTONIN GPROTEIN COUPLED TRANSMEMBRANE GLYCOPROTEIN MULTIGENE PD027821: M1-H84 G-PROTEIN COUPLED RECEPTORS DM00013 P31387 46-367: P46-T367 I48231 46-367: P46-T367 P47898 34-354: P46-T367 P20905 156-522: P46-S280	HMMER SPSCAN HMMER HMMER_Pfam BLIMPS_BLOCKS BLIMPS_PRINTS BLIMPS_PRINTS BLAST_PRODOM BLAST_DOMO

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
16	7480410CD1	324	S189 S194 S292 S314 S68	N9	Signal peptide: M1-A39 Transmembrane domains: G26-I50, C203-V217 7 transmembrane receptor (rhodopsin family): A42-Y291 G-protein coupled receptors signature: F103-V147 G-protein coupled receptor motif: G111-V127 G-protein coupled receptor BL00237: T283-A299, K91-P130, L208-Y219, R236-L262 Rhodopsin-like GPCR superfamily PR00237: L27-H51, M60-K81, Y105-V127, M200-L223, P25-L49, K273-A299 Olfactory receptor signature PR00245: M60-K81, F178-D192, F239-G254, V275-L286, S292-L306 G-protein coupled receptor: DM00013 P23275 17-306: H21-G307 A57069 15-304: H21-G307 P30954 29-316: L28-G307 S29709 11-299: P25-G307 Olfactory G-protein coupled receptor: PD000921: F169-M247 Olfactory G-protein coupled receptor: PD149621: V248-R308	SPScan HMMER HMMER-PFAM ProfileScan MOTIFS BLIMPS-BLOCKS BLIMPS-PRINTS BLIMPS-PRINTS BLAST-DOMO BLAST-PRODUM BLAST-PRODUM

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
17	55036418CD1	315	S194 S22 S292 S68 S8 S88	N6	Signal peptide: M1-A24 Transmembrane domains: L26-L46, C98-M119, F201-A220 7 transmembrane receptor (rhodopsin family): G42-Y291 G-protein coupled receptor motif: S111-I127 G-protein coupled receptors signature: L104-G147 G-protein coupled receptor BL00237: K91-P130, L208-Y219, K236-R262, T283-M299 Rhodopsin-like GPCR superfamily PR00237: V27-Y51, M60-K81, F105-I127, I200-L223, K273-M299 Olfactory receptor signature PR00245: M60-K81, F178-D192, L239-G254, A275-L286, S292-L306 G-protein coupled receptor: DM00013 P23275 17-306: I18-L302 A57069 15-304: F19-K304 P34982 17-305: V27-D307 P23270 18-311: L26-L306 Olfactory G-protein coupled receptor: PD000921: Y169-L246 Olfactory G-protein coupled receptor: PD149621: T247-R308	SPScan HMMER HMMER-PFAM MOTIFS ProfileScan BLIMPS-BLOCKS BLIMPS-PRINTS BLIMPS-PRINTS BLAST-DOMO BLAST-PRODOR BLAST-PRODOR

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
18	7481701CD1	324	S292 S314 S320 S67 S87 T136 T8		Transmembrane domains: L32-N52, V243-F263, Y267-N287 7 transmembrane receptor (rhodopsin family): G41-F180, V225-Y291 Visual pigments (opsins) retinal binding site: W271-T319 G-protein coupled receptor BL00237: R89-P128, R236-R262, S283-K299 Rhodopsin-like GPCR superfamily PR00237: V26-Q50, M59-K80, F103-I125, F12-F33, T144-F167, A238-R262, I273-K299 Olfactory receptor signature PR00245: M59-K80, I176-D190, F239-G254, S292-I306 Melanocortin receptor family PR00534: L51-I63, I125-T136 G-protein coupled receptor: DM00013 P23267 20-309: F17-I306 S29709 11-299: V29-G307 P23270 18-311: F17-K304 P23274 18-306: V26-I302 Olfactory G-protein coupled receptor: PD000921: L165-I247	HMMER HMMER-PFAM ProfileScan BLIMPS-BLOCKS BLIMPS-PRINTS BLIMPS-PRINTS BLIMPS-PRINTS BLAST-DOMO BLAST-PRODOM

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
19	7481774CD1	312	S16 S230 S264 S47 S65 T161 T289	N40	Transmembrane domains: I24-V46, Q98-M116, F198-L214 7 transmembrane receptor (rhodopsin family): G39-Y288 G-protein coupled receptor motif: V108-I124 G-protein coupled receptors signature: F100-T146 G-protein coupled receptor BL00237: K88-P127, V205-Y216, Q233-Q259, T280-K296 Olfactory receptor signature PR00245: M57-Q78, F175-S189, F236-G251, L272-L283, T289-L303 G-protein coupled receptor: DM00013 P23275 17-306: S16-L303 A57069 15-304: F15-L303 P34982 17-305: S16-L303 P30953 18-306: R18-L303 Olfactory G-protein coupled receptor: PD149621: T244-R307 Olfactory G-protein coupled receptor: PD000921: C167-L243	HMER HMER-PFAM MOTIFS ProfileScan BLIMPS-BLOCKS BLIMPS-PRINTS BLAST-DOMO BLAST-PRODOR BLAST-PRODOR

Table 4

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
20	7474806CB1	1076	1-1076	7075196H1 (BRAUTDR04) g7248967_edit g7407927_edit	1 533 1	532 1076 1102
21	7474840CB1	1102	1007-1102, 107-160, 486-522, 642-842			
22	7475092CB1	2529	628-1475, 1-81	55049853H1 71906055V1 1351856F6 (LATRTUT02) GNN.g7023955_000031_002 71900320V1 8017335J1 (BMATXE01) 55049805J1 7341260H1 (COLNDIN02) 70811587V1 6834479H1 (BRSTNON02) 70888652V1 7628842H1 (GBLADIE01) FL140044_00001	876 564 1757 1 1988 64 1256 307 1239 974 228 1 1	1612 1209 2367 468 2529 614 1956 991 1847 1575 660 284 2031
23	7341260CB1	1847	1-140, 1490-1847			
24	7473911CB1	2031	1-504, 669-1834			
25	7474767CB1	1130	1-1130	GNN.g6693326_000106_002 55093139J1 CpG.991027_B15_masked_fa. FL7475815_g8492585_000004_ g3892596	139 1 1 372	1130 276 723 1202
26	7475815CB1	1202	367-959, 1044-1202, 1-116			
27	60263275CB1	2079	134-490, 902-981	71704087V1 71651942V1 3642425T6 (LUNGNOT34) 524802R6 (CARCTXT01) 2435123H1 (BRAVUNT02) 71651560V1	85 986 1469 391 1 838	700 1727 2079 985 255 1544

Table 4

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
28	60203310CB1	5324	1-373, 963-3661, 4111-4162	71959831V1 60203311D1 7638002J1 (SEMTDE01) 491493H1 (HNT2AGT01) 71957696V1 4028716F7 (BRAINF023) 8103587H1 (MIXDDIE02) 9900324 8195081H2 (BRAIDIR04) 55073390J1 60200671D1 8195081J2 (BRAIDIR04) 55094091J1 55116120H1 70846216V1 GNN.g8140731 000028_002 GNN.g9309533 000030_002 2021568F6 (CONNOT01) 71243436V1 70844223V1 70845761V1 72398219V1 72374379V1 72373094V1 55049494J1 CPG_SAE300482544.R1 GNN:g6138786_000002_006.ed it 7290466F6 (BRAIFER06) 55049456J1 6925371H1 (PLACFER06) GNN:g9864547_000007_002.ed it 55084155J2 7341368F8 (COLNDIN02) g1507289	4179 3309 2283 3833 4646 4057 620 3652 1 1549 3444 432 1140 2658 1486 265 1 379 379 1272 872 804 675 971 814 1 1148 1 1576 461 249 472 1017 1 1879	5068 3622 2900 4073 5324 4787 1259 4165 623 2391 3823 1033 2026 3416 1962 1410 379 873 1931 1416 1415 1384 1558 1540 817 1481 1893 2304 1327 797 2103 1790 440 2322
29	7477349CB1	1962	1-1758			
30	55002225CB1	1558	80-1558, 1-49			
31	7475686CB1	2304	1-1212, 1548-2304			
32	7482007CB1	2322	1-628, 1452-1878, 724-1278			

Table 4

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
33	6769042CB1	2366	800-1248, 1-40, 1369-1717, 218-361, 1821-1861, 1930-2366	7157074H2 (BRAIFEJ02) 72138116D1 7629227H1 (GBLADIE01) 55147608J1	1169 1704 834 1	1797 2366 1341 908
34	7476053CB1	1458	1-1087	GNN.g7630808_000013_022 g3280262 GBI:g9454621.edit	1201 1088 1	1414 1458 1110
35	7480410CB1	975	1-816, 894- 975	55036194H1 (GPCRDPV02) GNN.g8979559_000011_002	166 1	371 975
36	55036418CB1	948	1-162, 193- 948	55036391H1 (GPCRDPV02) GNN.g7239420_000072_008	166 1	371 948
37	7481701CB1	1086	1-1086	GNN.g9795014_000002_002	1	1086
38	7481774CB1	1529	1-963	55143335J1 70822063V1 7361408F8 (BRAIFE05) 55142634H1	127 879 401 1	918 1529 1051 384

Table 5

Polynucleotide SEQ ID NO:	Incyte Project ID	Representative Library
20	7474806CB1	BRAUTDR04
22	7475092CB1	LATPTUT02
23	7341260CB1	COLNTUT03
24	7473911CB1	BRSTNOT23
27	60263275CB1	EOSITXT01
28	60203310CB1	BRAITUT01
29	7477349CB1	CONNNOT01
31	7475686CB1	BRAIFER06
32	7482007CB1	COLNDIN02
33	6769042CB1	GBLADIE01
38	7481774CB1	BRAIFEE05

Table 6

Library	Vector	Library Description
BRAIFEE05	PCDNA2.1	This 5' biased random primed library was constructed using RNA isolated from brain tissue removed from Caucasian male fetus who was still born with a hypoplastic left heart at 23 weeks' gestation. Serologies were negative.
BRAIFER06	PCDNA2.1	This random primed library was constructed using RNA isolated from brain tissue removed from a Caucasian male fetus who was stillborn with a hypoplastic left heart at 23 weeks' gestation. Serologies were negative.
BRAITUT01	PSPORT1	Library was constructed using RNA isolated from brain tumor tissue removed from a 50-year-old Caucasian female during a frontal lobectomy. Pathology indicated recurrent grade 3 oligoastrocytoma with focal necrosis and extensive calcification. Patient history included a speech disturbance and epilepsy. The patient's brain had also been irradiated with a total dose of 5,082 cGy (Fraction 8). Family history included a brain tumor.
BRAUTDR04	PCDNA2.1	This random primed library was constructed using RNA isolated from pooled striatum, dorsal caudate nucleus, dorsal putamen, and ventral nucleus accumbens tissue removed from a 55-year-old Caucasian female who died from cholangiocarcinoma. Pathology indicated mild meningeal fibrosis predominately over the convexities, scattered axonal spheroids in the white matter of the cingulate cortex and the thalamus, and a few scattered neurofibrillary tangles in the entorhinal cortex and the periaqueductal gray region. Pathology for the associated tumor tissue indicated well-differentiated cholangiocarcinoma of the liver with residual or relapsed tumor. Patient history included cholangiocarcinoma, post-operative Budd-Chiari syndrome, biliary ascites, hydrothorax, dehydration, malnutrition, oliguria and acute renal failure. Previous surgeries included cholecystectomy and resection of 85% of the liver.
BRSTNOT23	pINCY	Library was constructed using RNA isolated from diseased breast tissue removed from a 35-year-old Caucasian female during a bilateral reduction mammoplasty. Pathology indicated nonproliferative fibrocystic disease. Family history included type II diabetes, atherosclerotic coronary artery disease, acute myocardial infarction, hyperlipidemia, and coronary artery bypass.
COLNDIN02	pINCY	This normalized library was constructed from 4.72 million independent clones from a diseased colon and colon polyp tissue library. Starting RNA was made from pooled cDNA from two donors. cDNA was generated using mRNA isolated from diseased colon tissue removed from the cecum and descending colon of a 16-year-old Caucasian male (donor A) during partial colectomy, temporary ileostomy, and colonoscopy and from diseased colon polyp tissue removed from the cecum of a 67-year-old female (donor B). Pathology indicated innumerable (greater than 100) adenomatous polyps with low-grade dysplasia involving the entire colonic mucosa in the setting of familial polyposis coli (donor A), and a benign cecum polyp (donor B). Pathology for the associated tumor tissue (B) indicated invasive grade 3 adenocarcinoma that arose in tubulovillous adenoma forming a fungating mass in the cecum. The tumor infiltrated just through the muscularis propria. Multiple (2 of 17) regional lymph nodes were involved by metastatic adenocarcinoma. A tubulovillous adenoma and multiple (6) tubular adenomas with low-grade dysplasia were observed in the cecum and ascending colon. Donor A presented with abdominal pain and

Table 6 (cont.)

Library	Vector	Library Description
		Library Description flatulence. The patient was not taking any medications. Family history included benign colon neoplasm in the father and sibling(s); benign hypertension, cerebrovascular disease, breast cancer, uterine cancer, and type II diabetes in the grandparent(s).
COLNTUT03	pINCY	Library was constructed using RNA isolated from colon tumor tissue obtained from the sigmoid colon of a 62-year-old Caucasian male during a sigmoidectomy and permanent colostomy. Pathology indicated invasive grade 2 adenocarcinoma. One lymph node contained metastasis with extranodal extension. Patient history included hyperlipidemia, cataract disorder, and dermatitis. Family history included benign hypertension, atherosclerotic coronary artery disease, hyperlipidemia, breast cancer, and prostate cancer.
CONNOT01	pINCY	Library was constructed using RNA isolated from mesentery fat tissue obtained from a 71-year-old Caucasian male during a partial colectomy and permanent colostomy. Family history included atherosclerotic coronary artery disease, myocardial infarction, and extrinsic asthma.
EOSITXT01	pINCY	Library was constructed using RNA isolated from eosinophils stimulated with IL-5.
GBLADIE01	PCDNA2.1	This 5' biased random primed library was constructed using RNA isolated from diseased gallbladder tissue removed from a 55-year-old Caucasian female during laparoscopic cholecystectomy. Pathology indicated chronic cholecystitis and cholelithiasis (greater than 100 stones). The patient presented with cholelithiasis, abdominal pain, and tremors. Patient history included benign hypertension, Morton's neuroma, facial hirsutism, normal delivery, and tobacco abuse in remission. Previous surgeries included total abdominal hysterectomy, bilateral salpingo-oophorectomy, and adenotonsillectomy. Patient medications included Inderal and Premarin. Family history included breast cancer and ALS in the mother; chronic leukemia and ARDS in the father; breast cancer in the sibling(s); and atherosclerotic coronary artery disease in the grandparent(s).
LATRTUT02	pINCY	Library was constructed using RNA isolated from a myxoma removed from the left atrium of a 43-year-old Caucasian male during annuloplasty. Pathology indicated atrial myxoma. Patient history included pulmonary insufficiency, acute myocardial infarction, atherosclerotic coronary artery disease, hyperlipidemia, and tobacco use. Family history included benign hypertension, acute myocardial infarction, atherosclerotic coronary artery disease, and type II diabetes.

Table 7

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25:3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. USA 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183:63-98; and Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) Nucleic Acids Res. 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37:417-424.	Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol. 235:1501-1531; Sonnhammer, E.L.L. et al. (1998) Nucleic Acids Res. 26:320-322; Durbin, R. et al. (1998) Our World View, in a Nutshell, Cambridge Univ. Press, pp. 1-350.	PFAM hits: Probability value= 1.0E-3 or less Signal peptide hits: Score= 0 or greater

Table 7 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, M. et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score > GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score=3.5 or greater
TMAP	A program that uses weight matrices to delineate transmembrane segments on protein sequences and determine orientation.	Persson, B. and P. Argos (1994) J. Mol. Biol. 237:182-192; Persson, B. and P. Argos (1996) Protein Sci. 5:363-371.	
TMHMMER	A program that uses a hidden Markov model (HMM) to delineate transmembrane segments on protein sequences and determine orientation.	Sonnhammer, E.L. et al. (1998) Proc. Sixth Intl. Conf. on Intelligent Systems for Mol. Biol., Glasgow et al., eds., The Am. Assoc. for Artificial Intelligence Press, Menlo Park, CA, pp. 175-182.	
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide selected from the group consisting of:
 - a) a polypeptide comprising an amino acid sequence selected from the group consisting of
5 SEQ ID NO:1-19,
 - b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical
to an amino acid sequence selected from the group consisting of SEQ ID NO:1-19,
 - c) a biologically active fragment of a polypeptide having an amino acid sequence selected
from the group consisting of SEQ ID NO:1-19, and
 - 10 d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from
the group consisting of SEQ ID NO:1-19.
2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1-
19.
- 15 3. An isolated polynucleotide encoding a polypeptide of claim 1.
4. An isolated polynucleotide encoding a polypeptide of claim 2.
- 20 5. An isolated polynucleotide of claim 4 selected from the group consisting of SEQ ID
NO:20-38.
6. A recombinant polynucleotide comprising a promoter sequence operably linked to a
polynucleotide of claim 3.
- 25 7. A cell transformed with a recombinant polynucleotide of claim 6.
8. A transgenic organism comprising a recombinant polynucleotide of claim 6.
- 30 9. A method for producing a polypeptide of claim 1, the method comprising:
 - a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said
cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide
comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of
claim 1, and

b) recovering the polypeptide so expressed.

10. An isolated antibody which specifically binds to a polypeptide of claim 1.

5 11. An isolated polynucleotide selected from the group consisting of:

a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:20-38,

b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:20-38,

10 c) a polynucleotide complementary to a polynucleotide of a),

d) a polynucleotide complementary to a polynucleotide of b), and

e) an RNA equivalent of a)-d).

15 12. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 11.

13. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

20 a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and

b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

25

14. A method of claim 13, wherein the probe comprises at least 60 contiguous nucleotides.

15. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

30 a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and

b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

16. A composition comprising a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

17. A composition of claim 16, wherein the polypeptide has an amino acid sequence selected
5 from the group consisting of SEQ ID NO:1-19.

18. A method for treating a disease or condition associated with decreased expression of functional GCREC, comprising administering to a patient in need of such treatment the composition of claim 16.

10

19. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting agonist activity in the sample.

15

20. A composition comprising an agonist compound identified by a method of claim 19 and a pharmaceutically acceptable excipient.

21. A method for treating a disease or condition associated with decreased expression of
20 functional GCREC, comprising administering to a patient in need of such treatment a composition of claim 20.

22. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

25

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

23. A composition comprising an antagonist compound identified by a method of claim 22 and a pharmaceutically acceptable excipient.

30

24. A method for treating a disease or condition associated with overexpression of functional GCREC, comprising administering to a patient in need of such treatment a composition of claim 23.

25. A method of screening for a compound that specifically binds to the polypeptide of claim

1, said method comprising the steps of:

a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and

b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a
5 compound that specifically binds to the polypeptide of claim 1.

26. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, said method comprising:

a) combining the polypeptide of claim 1 with at least one test compound under conditions
10 permissive for the activity of the polypeptide of claim 1,

b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound,
and

c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change
15 in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

27. A method for screening a compound for effectiveness in altering expression of a target polynucleotide; wherein said target polynucleotide comprises a sequence of claim 5, the method
20 comprising:

a) exposing a sample comprising the target polynucleotide to a compound, under conditions suitable for the expression of the target polynucleotide,

b) detecting altered expression of the target polynucleotide, and

c) comparing the expression of the target polynucleotide in the presence of varying amounts
25 of the compound and in the absence of the compound.

28. A method for assessing toxicity of a test compound, said method comprising:

a) treating a biological sample containing nucleic acids with the test compound;

b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at
30 least 20 contiguous nucleotides of a polynucleotide of claim 11 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 11 or fragment thereof;

c) quantifying the amount of hybridization complex; and

d) comparing the amount of hybridization complex in the treated biological sample with the
35

amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

5 29. A diagnostic test for a condition or disease associated with the expression of GCREC in a biological sample comprising the steps of:

 a) combining the biological sample with an antibody of claim 10, under conditions suitable for the antibody to bind the polypeptide and form an antibody:polypeptide complex; and

 b) detecting the complex, wherein the presence of the complex correlates with the presence
10 of the polypeptide in the biological sample.

 30. The antibody of claim 10, wherein the antibody is:

 a) a chimeric antibody,

 b) a single chain antibody,

15 c) a Fab fragment,

 d) a F(ab')₂ fragment, or

 e) a humanized antibody.

 31. A composition comprising an antibody of claim 10 and an acceptable excipient.

20

 32. A method of diagnosing a condition or disease associated with the expression of GCREC in a subject, comprising administering to said subject an effective amount of the composition of claim 31.

25

 33. A composition of claim 31, wherein the antibody is labeled.

 34. A method of diagnosing a condition or disease associated with the expression of GCREC in a subject, comprising administering to said subject an effective amount of the composition of claim 33.

30

 35. A method of preparing a polyclonal antibody with the specificity of the antibody of claim 10 comprising:

 a) immunizing an animal with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, or an immunogenic fragment thereof, under conditions to

elicit an antibody response;

b) isolating antibodies from said animal; and

c) screening the isolated antibodies with the polypeptide, thereby identifying a polyclonal antibody which binds specifically to a polypeptide having an amino acid sequence selected from the

5 group consisting of SEQ ID NO:1-19.

36. An antibody produced by a method of claim 35.

37. A composition comprising the antibody of claim 36 and a suitable carrier.

10

38. A method of making a monoclonal antibody with the specificity of the antibody of claim 10 comprising:

a) immunizing an animal with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19, or an immunogenic fragment thereof, under conditions to

15 elicit an antibody response;

b) isolating antibody producing cells from the animal;

c) fusing the antibody producing cells with immortalized cells to form monoclonal antibody-producing hybridoma cells;

d) culturing the hybridoma cells; and

20 e) isolating from the culture monoclonal antibody which binds specifically to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19.

39. A monoclonal antibody produced by a method of claim 38.

25

40. A composition comprising the antibody of claim 39 and a suitable carrier.

41. The antibody of claim 10, wherein the antibody is produced by screening a Fab expression library.

30

42. The antibody of claim 10, wherein the antibody is produced by screening a recombinant immunoglobulin library.

43. A method for detecting a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19 in a sample, comprising the steps of:

- a) incubating the antibody of claim 10 with a sample under conditions to allow specific binding of the antibody and the polypeptide; and
- b) detecting specific binding, wherein specific binding indicates the presence of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19 in the sample.

44. A method of purifying a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19 from a sample, the method comprising:

- a) incubating the antibody of claim 10 with a sample under conditions to allow specific binding of the antibody and the polypeptide; and
- b) separating the antibody from the sample and obtaining the purified polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19.

45. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:1.

46. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:2.

47. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:3.

48. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:4.

49. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:5.

50. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:6.

51. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:7.

52. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:8.

53. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:9.

54. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:10.

55. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:11.

56. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:12.
57. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:13.
- 5 58. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:14.
59. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:15.
60. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:16.
- 10 61. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:17.
62. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:18.
- 15 63. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:19.
64. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:20.
- 20 65. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:21.
66. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:22.
- 25 67. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:23.
68. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
30 NO:24.
69. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:25.

70. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:26.

71. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
5 NO:27.

72. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:28.

10 73. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:29.

74. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:30.
15

75. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:31.

76. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
20 NO:32.

77. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:33.

25 78. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:34.

79. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:35.
30

80. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:36.

81. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID

NO:37.

82. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:38.

<110> INCYTE GENOMICS, INC.

THORNTON, Michael

PATERSON, Chandra

LAL, Preeti

BURFORD, Neil

YUE, Henry

GANDHI, Aameena R.

ELLIOTT, Vicki S.

RAMKUMAR, Jayalaxmi

BAUGHN, Mariah R.

KALLICK, Deborah A.

WALIA, Narinder K.

HAFALIA, April J.A.

YAO, Monique G.

LU, Yan

TRIBOULEY, Catherine M.

POLICKY, Jennifer L.

KEARNEY, Liam

GRAUL, Richard

WARREN, Bridget

LEE, Ernestine A.

DING, Li

<120> G-PROTEIN COUPLED RECEPTORS

<130> PI-0176 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/221,478; 60/223,268; 60/227,054; 60/231,121; 60/232,243;
60/232,691; 60/235,146151> 2000-07-27; 2000-08-03; 2000-08-21; 2000-09-08; 2000-09-13;
2000-09-15; 2000-09-22

<160> 38

<170> PERL Program

<210> 1

<211> 339

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7474806CD1

<400> 1

Met	Leu	Ser	Ile	Leu	Leu	Pro	Ser	Arg	Gly	Ser	Arg	Ser	Gly	Ser
1				5					10					15
Arg	Arg	Gly	Ala	Leu	Leu	Glu	Gly	Ala	Ser	Arg	Asp	Met	Glu	
				20					25					30
Lys	Val	Asp	Met	Asn	Thr	Ser	Gln	Glu	Gln	Gly	Leu	Cys	Gln	Phe
				35					40					45
Ser	Glu	Lys	Tyr	Lys	Gln	Val	Tyr	Leu	Ser	Leu	Ala	Tyr	Ser	Ile
				50					55					60
Ile	Phe	Ile	Leu	Gly	Leu	Pro	Leu	Asn	Gly	Thr	Val	Leu	Trp	His
				65					70					75
Ser	Trp	Gly	Gln	Thr	Lys	Arg	Trp	Ser	Cys	Ala	Thr	Thr	Tyr	Leu
				80					85					90
Val	Asn	Leu	Met	Val	Ala	Asp	Leu	Leu	Tyr	Val	Leu	Leu	Pro	Phe
				95					100					105
Leu	Ile	Ile	Thr	Tyr	Ser	Leu	Asp	Asp	Arg	Trp	Pro	Phe	Gly	Glu

	110		115		120
Leu Leu Cys Lys	Leu Val His Phe Leu	Phe Tyr Ile Asn Leu Tyr			
	125		130		135
Gly Ser Ile Leu	Leu Leu Thr Cys Ile	Ser Val His Gln Phe Leu			
	140		145		150
Gly Val Trp His	Pro Leu Cys Ser Leu	Pro Tyr Arg Thr Arg Arg			
	155		160		165
His Ala Trp Leu	Gly Thr Ser Thr Thr	Trp Ala Leu Val Val Leu			
	170		175		180
Gln Leu Leu Pro	Thr Leu Ala Phe Ser	His Thr Asp Tyr Ile Asn			
	185		190		195
Gly Gln Met Ile	Trp Tyr Asp Met Thr	Ser Gln Glu Asn Phe Asp			
	200		205		210
Arg Leu Phe Ala	Tyr Gly Ile Val Leu	Thr Leu Ser Gly Phe Leu			
	215		220		225
Ser Pro Ser Leu	Val Ile Leu Val Cys	Tyr Ser Leu Met Val Arg			
	230		235		240
Ser Leu Ile Lys	Pro Glu Glu Asn Leu	Met Arg Thr Gly Asn Thr			
	245		250		255
Ala Arg Ala Arg	Ser Ile Arg Thr Ile	Leu Leu Val Cys Gly Leu			
	260		265		270
Phe Thr Leu Cys	Phe Val Pro Phe His	Ile Thr Arg Ser Phe Tyr			
	275		280		285
Leu Thr Ile Cys	Phe Leu Leu Ser Gln	Asp Cys Gln Leu Leu Met			
	290		295		300
Ala Pro Ser Val	Ala Tyr Lys Ile Trp	Arg Pro Leu Val Ser Val			
	305		310		315
Ser Ser Cys Leu	Asn Pro Val Leu Tyr	Phe Leu Ser Arg Gly Ala			
	320		325		330
Lys Ile Glu Ser	Gly Ser Ser Arg Asn				
	335				

<210> 2

<211> 335

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7474840CD1

<400> 2

Met Thr Pro Gly Gly	Arg Ala Cys Ser	Glu Met Arg Ser Cys His
1	5	10 15
Cys Ala Pro Ala Trp	Ala Thr Glu Arg Asp	Ser Val Ser Lys Lys
20	25	30
Lys Lys Asn Lys Lys	Lys Asn Leu Phe Ser	Gln Ala Thr Ile Gly
35	40	45
Leu Leu Ala Asn Thr	Phe Phe Leu Phe Phe	Asn Ile Phe Ile Phe
50	55	60
Leu Gln Asp Gln Lys	Ser Lys Pro His Asp	Leu Ile Ser Cys Asn
65	70	75
Ser Ala Phe Ile His	Val Val Met Phe Leu	Thr Val Val Asp Ala
80	85	90
Trp Pro Pro Asp Met	Pro Glu Ser Leu His	Leu Gly Asn Glu Phe
95	100	105
Lys Phe Lys Ser Leu	Ser Tyr Ile Asn Arg	Val Arg Met Gly Leu
110	115	120
Cys Ile Cys Asn Ile	Cys Leu Leu Ser Ile	His Gln Ala Asn Thr
125	130	135
Ile Ser Pro Asn Asn	Phe Cys Leu Ala Arg	Leu Lys Gln Lys Phe
140	145	150
Thr Asn Asn Ile Ile	Met Ser Ser Phe Phe	Ser Phe Phe Phe Trp

Ser Ile Asn Leu Ser Phe Ser Tyr Asn Ile Val Phe Phe Thr Val	155	160	165
	170	175	180
Ala Ser Ser Asn Val Thr Gln Asn Ser Leu Pro Lys Gly Ser Asn	185	190	195
Thr Val His Phe Leu Pro Met Lys Ser Phe Met Arg Lys Val Phe	200	205	210
Phe Thr Leu Thr Leu Ser Arg Asp Val Phe Ile Ile Gly Ile Thr	215	220	225
Leu His Ser Ile Ala His Met Val Ile Leu Val Ser Arg His Glu	230	235	240
Thr Gln Ser Gln His Leu His Ser Ile Ser Ile Ser Pro Gln Ala	245	250	255
Phe Pro Glu Lys Arg Ala Ala Gln Thr Ile Pro Leu Leu Val Ser	260	265	270
Tyr Cys Leu Val Met Cys Trp Val Asp Leu Ile Ile Ser Ser Ser	275	280	285
Ser Thr Leu Leu Trp Thr Cys Asn Pro Val Phe Leu Ser Met Gln	290	295	300
Asn Leu Val Gly Asp Val Tyr Ala Thr Val Val Leu Leu Glu Gln	305	310	315
Ile Ser Ser Asp Lys Asn Ile Val Asp Ile Leu Gln Asn Met Gln	320	325	330
Ser Ala Ile Lys Leu	335		

<210> 3

<211> 428

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7475092CD1

<400> 3

Met Gln Arg Lys Glu Lys Ala Lys Cys Pro Gln Glu Ala Pro Ala	1	5	10	15
Gly Arg Glu Pro Ser Thr Pro Gly Gly Gly Ser Gly Gly Gly Gly	20	25	30	35
Ala Val Ala Ala Ala Ser Gly Ala Ala Val Pro Gly Ser Val Gln	40	45	50	55
Leu Ala Leu Ser Val Leu His Ala Leu Leu Tyr Ala Ala Leu Phe	60	65	70	75
Ala Phe Ala Tyr Leu Gln Leu Trp Arg Leu Leu Leu Tyr Arg Glu	80	85	90	95
Arg Arg Leu Ser Tyr Gln Ser Leu Cys Leu Phe Leu Cys Leu Leu	100	105	110	115
Trp Ala Ala Leu Arg Thr Thr Leu Phe Ser Ala Ala Phe Ser Leu	120	125	130	135
Ser Gly Ser Leu Pro Leu Leu Arg Pro Pro Ala His Leu His Phe	140	145	150	155
Phe Pro His Trp Leu Leu Tyr Cys Phe Pro Ser Cys Leu Gln Phe	160	165	170	175
Ser Thr Leu Cys Leu Leu Asn Leu Tyr Leu Ala Glu Val Ile Cys	180	185	190	195
Lys Val Arg Cys Ala Thr Glu Leu Asp Arg His Lys Ile Leu Leu	200	205	210	215
His Leu Gly Phe Ile Met Ala Ser Leu Leu Phe Leu Val Val Asn	220	225	230	235
Leu Thr Cys Ala Met Leu Val His Gly Asp Val Pro Glu Asn Gln	240	245	250	255
Leu Lys Trp Thr Val Phe Val Arg Ala Leu Ile Asn Asp Ser Leu	260	265	270	275

Phe Ile Leu Cys	200	205	210
Ala Ile Ser Leu Val		Cys Tyr Ile Cys Lys	Ile
215		220	225
Thr Lys Met Ser	Ser Ala Asn Val Tyr	Leu Glu Ser Lys Gly	Met
230		235	240
Ser Leu Cys Gln	Thr Val Val Val Gly	Ser Val Val Ile Leu	Leu
245		250	255
Tyr Ser Ser Arg	Ala Cys Tyr Asn Leu	Val Val Val Thr Ile	Ser
260		265	270
Gln Asp Thr Leu	Glu Ser Pro Phe Asn Tyr	Gly Trp Asp Asn	Leu
275		280	285
Ser Asp Lys Ala	His Val Glu Asp Ile	Ser Gly Glu Glu Tyr	Ile
290		295	300
Val Phe Gly Met	Val Leu Phe Leu Trp	Glu His Val Pro Ala	Trp
305		310	315
Ser Val Val Leu	Phe Phe Arg Ala Gln	Arg Leu Asn Gln Asn	Leu
320		325	330
Ala Pro Ala Gly	Met Ile Asn Ser His	Ser Tyr Ser Ser Arg	Ala
335		340	345
Tyr Phe Phe Asp	Asn Pro Arg Arg Tyr	Asp Ser Asp Asp Asp	Leu
350		355	360
Pro Arg Leu Gly	Ser Ser Arg Glu Gly	Ser Leu Pro Asn Ser	Gln
365		370	375
Ser Leu Gly Trp	Tyr Gly Thr Met Thr	Gly Cys Gly Ser Ser	Ser
380		385	390
Tyr Thr Val Thr	Pro His Leu Asn Gly	Pro Met Thr Asp Thr	Ala
395		400	405
Pro Leu Leu Phe	Thr Cys Ser Asn Leu	Asp Leu Asn Asn His	His
410		415	420
Ser Leu Tyr Val	Thr Pro Gln Asn		
425			

<210> 4

<211> 330

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7341260CD1

<400> 4

Met Thr Pro Asn Ser	Thr Gly Glu Val	Pro Ser Pro Ile Pro	Lys
1	5	10	15
Gly Ala Leu Gly Leu	Ser Leu Ala Leu	Ala Ser Leu Ile Ile	Thr
20	25		30
Ala Asn Leu Leu Leu	Ala Leu Gly Ile	Ala Trp Asp Arg Arg	Leu
35	40		45
Arg Ser Pro Pro Ala	Gly Cys Phe Phe	Leu Ser Leu Leu Leu	Ala
50	55		60
Gly Leu Leu Thr Gly	Leu Ala Leu Pro	Thr Leu Pro Gly Leu	Trp
65	70		75
Asn Gln Ser Arg Arg	Gly Tyr Trp Ser	Cys Leu Leu Val Tyr	Leu
80	85		90
Ala Pro Asn Phe Ser	Phe Leu Ser Leu	Leu Ala Asn Leu Leu	Leu
95	100		105
Val His Gly Glu Arg	Tyr Met Ala Val	Leu Arg Pro Leu Gln	Pro
110	115		120
Pro Gly Ser Ile Arg	Leu Ala Leu Leu	Leu Thr Trp Ala Gly	Pro
125	130		135
Leu Leu Phe Ala Ser	Leu Pro Ala Leu	Gly Trp Asn His Trp	Thr
140	145		150
Pro Gly Ala Asn Cys	Ser Ser Gln Ala	Ile Phe Pro Ala Pro	Tyr

155	160	165
Leu Tyr Leu Glu Val Tyr Gly Leu Leu	Leu Pro Ala Val Gly Ala	
170	175	180
Ala Ala Phe Leu Ser Val Arg Val Leu	Ala Thr Ala His Arg Gln	
185	190	195
Leu Gln Asp Ile Cys Arg Leu Glu Arg	Ala Val Cys Arg Asp Glu	
200	205	210
Pro Ser Ala Leu Ala Arg Ala Leu Thr	Trp Arg Gln Ala Arg Ala	
215	220	225
Gln Ala Gly Ala Met Leu Leu Phe Gly	Leu Cys Trp Gly Pro Tyr	
230	235	240
Val Ala Thr Leu Leu Leu Ser Val Leu	Ala Tyr Glu Gln Arg Pro	
245	250	255
Pro Leu Gly Pro Gly Thr Leu Leu Ser	Leu Leu Ser Leu Gly Ser	
260	265	270
Ala Ser Ala Ala Ala Val Pro Val Ala	Met Gly Leu Gly Asp Gln	
275	280	285
Arg Tyr Thr Ala Pro Trp Arg Ala Ala	Ala Gln Arg Cys Leu Gln	
290	295	300
Gly Leu Trp Gly Arg Ala Ser Arg Asp	Ser Pro Gly Pro Ser Ile	
305	310	315
Ala Tyr His Pro Ser Ser Gln Ser Ser	Val Asp Leu Asp Leu Asn	
320	325	330

<210> 5

<211> 676

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7473911CD1

<400> 5

Met Asn Lys Asn Asn Lys Pro Ser Ser Phe Ile Ala Ile Arg Asn	
1 5 10 15	
Ala Ala Phe Ser Glu Val Gly Ile Gly Ile Ser Ala Asn Ala Met	
20 25 30	
Leu Leu Leu Phe His Ile Leu Thr Cys Leu Leu Lys His Arg Thr	
35 40 45	
Lys Pro Ala Asp Leu Ile Val Cys His Val Ala Leu Ile His Ile	
50 55 60	
Ile Leu Leu Leu Pro Thr Glu Phe Ile Ala Thr Asp Ile Phe Gly	
65 70 75	
Ser Gln Asp Ser Glu Asp Asp Ile Lys His Lys Ser Val Ile Tyr	
80 85 90	
Arg Arg Asn Arg Gln Ser Gln His Phe His Ser Thr Asn Leu Ser	
95 100 105	
Pro Lys Ala Pro Pro Glu Lys Met Ala Thr Gln Thr Ile Leu Leu	
110 115 120	
Leu Val Ser Cys Phe Val Ile Val Tyr Val Leu Asp Cys Val Val	
125 130 135	
Ala Ser Cys Ser Gly Leu Val Trp Asn Ser Asp Pro Val Arg His	
140 145 150	
Arg Val Gln Met Leu Val Asp Asn Gly Tyr Ala Thr Ile Ser Pro	
155 160 165	
Ser Val Leu Pro Arg Leu Thr Ala Pro Asn Glu Trp Arg Ala Ser	
170 175 180	
Val Tyr Leu Asn Asp Ser Leu Asn Lys Cys Ser Asn Gly Arg Leu	
185 190 195	
Leu Cys Val Asp Arg Gly Leu Asp Glu Gly Pro Arg Ser Val Pro	
200 205 210	

Lys	Cys	Ser	Glu	Ser	Glu	Thr	Asp	Glu	Asp	Tyr	Ile	Val	Leu	Arg
				215					220					225
Ala	Pro	Leu	Arg	Glu	Asp	Glu	Pro	Lys	Asp	Gly	Gly	Ser	Val	Gly
				230					235					240
Asn	Ala	Ala	Leu	Val	Ser	Pro	Glu	Ala	Ser	Ala	Glu	Glu	Glu	Glu
				245					250					255
Glu	Arg	Glu	Glu	Gly	Gly	Glu	Ala	Cys	Gly	Leu	Glu	Arg	Thr	Gly
				260					265					270
Ala	Gly	Gly	Glu	Gln	Val	Asp	Leu	Gly	Glu	Leu	Pro	Asp	His	Glu
				275					280					285
Glu	Lys	Ser	Asn	Gln	Lys	Val	Ala	Ala	Ala	Thr	Leu	Glu	Asp	Arg
				290					295					300
Thr	Gln	Asp	Glu	Pro	Ala	Glu	Glu	Ser	Cys	Gln	Ile	Val	Leu	Phe
				305					310					315
Gln	Asn	Asn	Cys	Met	Asp	Asn	Phe	Val	Thr	Ser	Leu	Thr	Gly	Ser
				320					325					330
Pro	Tyr	Glu	Phe	Phe	Pro	Thr	Lys	Ser	Thr	Ser	Phe	Cys	Arg	Glu
				335					340					345
Ser	Cys	Ser	Pro	Phe	Ser	Glu	Ser	Val	Lys	Ser	Leu	Glu	Ser	Glu
				350					355					360
Gln	Ala	Pro	Lys	Leu	Gly	Leu	Cys	Ala	Glu	Glu	Asp	Pro	Val	Val
				365					370					375
Gly	Ala	Leu	Cys	Gly	Gln	His	Gly	Pro	Leu	Gln	Asp	Gly	Val	Ala
				380					385					390
Glu	Gly	Pro	Thr	Ala	Pro	Asp	Val	Val	Val	Leu	Pro	Lys	Glu	Glu
				395					400					405
Glu	Lys	Glu	Glu	Val	Ile	Val	Asp	Asp	Met	Leu	Ala	Asn	Pro	Tyr
				410					415					420
Val	Met	Gly	Asp	Glu	Gly	Glu	Glu	Glu	Glu	Glu	Glu	Phe	Val	Asp
				425					430					435
Asp	Thr	Leu	Ala	Asn	Pro	Tyr	Val	Met	Gly	Val	Gly	Leu	Pro	Gly
				440					445					450
Arg	Gly	Gly	Glu	Glu	Glu	Glu	Glu	Glu	Glu	Val	Val	Asp	Asp	Thr
				455					460					465
Leu	Ala	Ser	Leu	Tyr	Lys	Met	Gly	Glu	Glu	His	Arg	His	Lys	Gly
				470					475					480
Leu	Ala	Pro	Leu	Trp	Glu	Gly	Gly	Gln	Lys	Pro	Ser	Gln	Lys	Leu
				485					490					495
Pro	Pro	Lys	Lys	Pro	Asp	Leu	Arg	Gln	Val	Pro	Gln	Pro	Leu	Ala
				500					505					510
Ser	Glu	Val	Pro	Gln	Arg	Arg	Gln	Glu	Arg	Ala	Val	Val	Thr	Glu
				515					520					525
Gly	Arg	Pro	Leu	Glu	Ala	Ser	Arg	Ala	Leu	Pro	Ala	Lys	Pro	Arg
				530					535					540
Ala	Phe	Thr	Leu	Tyr	Pro	Arg	Ser	Phe	Ser	Val	Glu	Gly	Gln	Glu
				545					550					555
Ile	Pro	Val	Ser	Ile	Ser	Val	Tyr	Trp	Glu	Pro	Glu	Gly	Ser	Gly
				560					565					570
Leu	Asp	Asp	His	Arg	Ile	Lys	Arg	Lys	Glu	Glu	His	Leu	Ser	Val
				575					580					585
Val	Ser	Gly	Ser	Phe	Ser	Gln	Arg	Asn	His	Leu	Pro	Ser	Ser	Gly
				590					595					600
Thr	Ser	Thr	Pro	Ser	Ser	Met	Val	Asp	Ile	Pro	Pro	Pro	Phe	Asp
				605					610					615
Leu	Ala	Cys	Ile	Thr	Lys	Lys	Pro	Ile	Thr	Lys	Ser	Ser	Pro	Ser
				620					625					630
Leu	Leu	Ile	Asp	Ser	Asp	Ser	Pro	Asp	Lys	Tyr	Lys	Lys	Lys	Lys
				635					640					645
Ser	Ser	Phe	Lys	Arg	Phe	Leu	Ala	Leu	Met	Phe	Asn	Lys	Met	Glu
				650					655					660
Arg	Pro	Gly	Thr	Met	Ala	His	Ala	Cys	His	Pro	Ser	Thr	Leu	Gly
				665					670					675
Ser														

<210> 6
 <211> 372
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7474767CD1

<400> 6
 Met Glu His Thr His Ala His Leu Ala Ala Asn Ser Ser Leu Ser
 1 5 10 15
 Trp Trp Ser Pro Gly Ser Ala Cys Gly Leu Gly Phe Val Pro Val
 20 25 30
 Val Tyr Tyr Ser Leu Leu Leu Cys Leu Gly Leu Pro Ala Asn Ile
 35 40 45
 Leu Thr Val Ile Ile Leu Ser Gln Leu Val Ala Arg Arg Gln Lys
 50 55 60
 Ser Ser Tyr Asn Tyr Leu Leu Ala Leu Ala Ala Ala Asp Ile Leu
 65 70 75
 Val Leu Phe Phe Ile Val Phe Val Asp Phe Leu Leu Glu Asp Phe
 80 85 90
 Ile Leu Asn Met Gln Met Pro Gln Val Pro Asp Lys Ile Ile Glu
 95 100 105
 Val Leu Glu Phe Ser Ser Ile His Thr Ser Ile Trp Ile Thr Val
 110 115 120
 Pro Leu Thr Ile Asp Arg Tyr Ile Ala Val Cys His Pro Leu Lys
 125 130 135
 Tyr His Thr Val Ser Tyr Pro Ala Arg Thr Arg Lys Val Ile Val
 140 145 150
 Ser Val Tyr Ile Thr Cys Phe Leu Thr Ser Ile Pro Tyr Tyr Trp
 155 160 165
 Trp Pro Asn Ile Trp Thr Glu Asp Tyr Ile Ser Thr Ser Val His
 170 175 180
 His Val Leu Ile Trp Ile His Cys Phe Thr Val Tyr Leu Val Pro
 185 190 195
 Cys Ser Ile Phe Phe Ile Leu Asn Ser Ile Ile Val Tyr Lys Leu
 200 205 210
 Arg Arg Lys Ser Asn Phe Arg Leu Arg Gly Tyr Ser Thr Gly Lys
 215 220 225
 Thr Thr Ala Ile Leu Phe Thr Ile Thr Ser Ile Phe Ala Thr Leu
 230 235 240
 Trp Ala Pro Arg Ile Ile Met Ile Leu Tyr His Leu Tyr Gly Ala
 245 250 255
 Pro Ile Gln Asn Arg Trp Leu Val His Ile Met Ser Asp Ile Ala
 260 265 270
 Asn Met Leu Ala Leu Leu Asn Thr Ala Ile Asn Phe Phe Leu Tyr
 275 280 285
 Cys Phe Ile Ser Lys Arg Phe Arg Thr Met Ala Ala Ala Thr Leu
 290 295 300
 Lys Ala Phe Phe Lys Cys Gln Lys Gln Pro Val Gln Phe Tyr Thr
 305 310 315
 Asn His Asn Phe Ser Ile Thr Ser Ser Pro Trp Ile Ser Pro Ala
 320 325 330
 Asn Ser His Cys Ile Lys Met Leu Val Tyr Gln Tyr Asp Lys Asn
 335 340 345
 Gly Lys Pro Ile Lys Ser Arg Asn Asp Ser Lys Ser Ser Tyr Gln
 350 355 360
 Phe Glu Asp Ala Ile Gly Ala Cys Val Ile Ile Leu
 365 370

<210> 7
 <211> 271
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7475815CD1

<400> 7
 Met Asn Lys Asn Asn Lys Pro Ser Ser Phe Ile Ala Ile Arg Asn
 1 5 10 15
 Ala Ala Phe Ser Glu Val Gly Ile Gly Ile Ser Ala Asn Ala Met
 20 25 30
 Leu Leu Leu Phe His Ile Leu Thr Cys Leu Leu Lys His Arg Thr
 35 40 45
 Lys Pro Ala Asp Leu Ile Val Cys His Val Ala Leu Ile His Ile
 50 55 60
 Ile Leu Leu Leu Pro Thr Glu Phe Ile Ala Thr Asp Ile Phe Gly
 65 70 75
 Ser Gln Asp Ser Glu Asp Asp Ile Lys His Lys Ser Val Ile Tyr
 80 85 90
 Arg Tyr Arg Leu Met Arg Gly Leu Ser Ile Ser Thr Thr Cys Leu
 95 100 105
 Leu Ser Ile Leu Pro Ala Ile Thr Cys Ser Pro Arg Ser Ser Cys
 110 115 120
 Leu Ala Val Phe Lys Asp Ser His Ile Thr Asn His Val Ala Phe
 125 130 135
 Ser Ser Val Phe His Ile Ser Ile Ser Asp Ser Phe Leu Val Ser
 140 145 150
 Thr Leu Pro Ile Lys Asn Leu Ala Ser Asn Ser Leu Thr Phe Val
 155 160 165
 Thr Gln Ser Cys Ser Ala Gly Ile Gly Ser Arg Pro Pro Ser Ser
 170 175 180
 Gly Tyr Met Val Ile Leu Leu Ser Arg Arg Asn Arg Gln Ser Gln
 185 190 195
 His Phe His Ser Thr Asn Leu Ser Pro Lys Ala Pro Pro Glu Lys
 200 205 210
 Met Ala Thr Gln Thr Ile Leu Leu Leu Val Ser Cys Phe Val Ile
 215 220 225
 Val Tyr Val Leu Asp Cys Val Val Ala Ser Cys Ser Gly Leu Val
 230 235 240
 Trp Asn Ser Asp Pro Val Arg His Arg Val Gln Met Leu Val Asp
 245 250 255
 Asn Gly Tyr Ala Thr Ile Ser Pro Ser Val Leu Val Ser Thr Glu
 260 265 270
 Lys

<210> 8
 <211> 611
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 60263275CD1

<400> 8
 Met Gln Gly Pro Leu Leu Leu Pro Gly Leu Cys Phe Leu Leu Ser
 1 5 10 15
 Leu Phe Gly Ala Val Thr Gln Lys Thr Lys Asn Ile Asn Glu Cys
 20 25 30

Thr	Pro	Pro	Tyr	Ser	Val	Tyr	Cys	Gly	Phe	Asn	Ala	Val	Cys	Tyr
				35					40					45
Asn	Val	Glu	Gly	Ser	Phe	Tyr	Cys	Gln	Cys	Val	Pro	Gly	Tyr	Arg
				50					55					60
Leu	His	Ser	Gly	Asn	Glu	Gln	Phe	Ser	Asn	Ser	Asn	Glu	Asn	Thr
				65					70					75
Cys	Gln	Asp	Thr	Thr	Ser	Ser	Lys	Thr	Thr	Gln	Gly	Arg	Lys	Glu
				80					85					90
Leu	Gln	Lys	Ile	Val	Asp	Lys	Phe	Glu	Ser	Leu	Leu	Thr	Asn	Gln
				95					100					105
Thr	Leu	Trp	Arg	Thr	Glu	Gly	Arg	Gln	Glu	Ile	Ser	Ser	Thr	Ala
				110					115					120
Thr	Thr	Ile	Leu	Arg	Asp	Val	Glu	Ser	Lys	Val	Leu	Glu	Thr	Ala
				125					130					135
Leu	Lys	Asp	Pro	Glu	Gln	Lys	Val	Leu	Lys	Ile	Gln	Asn	Asp	Ser
				140					145					150
Val	Ala	Ile	Glu	Thr	Gln	Ala	Ile	Thr	Asp	Asn	Cys	Ser	Glu	Glu
				155					160					165
Arg	Lys	Thr	Phe	Asn	Leu	Asn	Val	Gln	Met	Asn	Ser	Met	Asp	Ile
				170					175					180
Arg	Cys	Ser	Asp	Ile	Ile	Gln	Gly	Asp	Thr	Gln	Gly	Pro	Ser	Ala
				185					190					195
Ile	Ala	Phe	Ile	Ser	Tyr	Ser	Ser	Leu	Gly	Asn	Ile	Ile	Asn	Ala
				200					205					210
Thr	Phe	Phe	Glu	Glu	Met	Asp	Lys	Lys	Asp	Gln	Val	Tyr	Leu	Asn
				215					220					225
Ser	Gln	Val	Val	Ser	Ala	Ala	Ile	Gly	Pro	Lys	Arg	Asn	Val	Ser
				230					235					240
Leu	Ser	Lys	Ser	Val	Thr	Leu	Thr	Phe	Gln	His	Val	Lys	Met	Thr
				245					250					255
Pro	Ser	Thr	Lys	Lys	Val	Phe	Cys	Val	Tyr	Trp	Lys	Ser	Thr	Gly
				260					265					270
Gln	Gly	Ser	Gln	Trp	Ser	Arg	Asp	Gly	Cys	Phe	Leu	Ile	His	Val
				275					280					285
Asn	Lys	Ser	His	Thr	Met	Cys	Asn	Cys	Ser	His	Leu	Ser	Ser	Phe
				290					295					300
Ala	Val	Leu	Met	Ala	Leu	Thr	Ser	Gln	Glu	Glu	Asp	Pro	Val	Leu
				305					310					315
Thr	Val	Ile	Thr	Tyr	Val	Gly	Leu	Ser	Val	Ser	Leu	Leu	Cys	Leu
				320					325					330
Leu	Leu	Ala	Ala	Leu	Thr	Phe	Leu	Leu	Cys	Lys	Ala	Ile	Gln	Asn
				335					340					345
Thr	Ser	Thr	Ser	Leu	His	Leu	Gln	Leu	Ser	Leu	Cys	Leu	Phe	Leu
				350					355					360
Ala	His	Leu	Leu	Phe	Leu	Val	Gly	Ile	Asp	Arg	Thr	Glu	Pro	Lys
				365					370					375
Val	Leu	Cys	Ser	Ile	Ile	Ala	Gly	Ala	Leu	His	Tyr	Leu	Tyr	Leu
				380					385					390
Ala	Ala	Phe	Thr	Trp	Met	Leu	Leu	Glu	Gly	Val	His	Leu	Phe	Leu
				395					400					405
Thr	Ala	Arg	Asn	Leu	Thr	Val	Val	Asn	Tyr	Ser	Ser	Ile	Asn	Arg
				410					415					420
Leu	Met	Lys	Trp	Ile	Met	Phe	Pro	Val	Gly	Tyr	Gly	Val	Pro	Ala
				425					430					435
Val	Thr	Val	Ala	Ile	Ser	Ala	Ala	Ser	Trp	Pro	His	Leu	Tyr	Gly
				440					445					450
Thr	Ala	Asp	Arg	Cys	Trp	Leu	His	Leu	Asp	Gln	Gly	Phe	Met	Trp
				455					460					465
Ser	Phe	Leu	Gly	Pro	Val	Cys	Ala	Ile	Phe	Ser	Ala	Asn	Leu	Val
				470					475					480
Leu	Phe	Ile	Leu	Val	Phe	Trp	Ile	Leu	Lys	Arg	Lys	Leu	Ser	Ser
				485					490					495
Leu	Asn	Ser	Glu	Val	Ser	Thr	Ile	Gln	Asn	Thr	Arg	Met	Leu	Ala

	500		505		510
Phe Lys Ala Thr	Ala Gln Leu Phe Ile	Leu Gly Cys Thr Trp Cys			
	515	520			525
Leu Gly Leu Leu	Gln Val Gly Pro Ala	Ala Gln Val Met Ala Tyr			
	530	535			540
Leu Phe Thr Ile	Ile Asn Ser Leu Gln	Gly Phe Phe Ile Phe			Leu
	545	550			555
Val Tyr Cys Leu	Leu Ser Gln Gln Val	Gln Lys Gln Tyr Gln			Lys
	560	565			570
Trp Phe Arg Glu	Ile Val Lys Ser Lys	Ser Glu Ser Glu Thr Tyr			
	575	580			585
Thr Leu Ser Ser	Lys Met Gly Pro Asp	Ser Lys Pro Ser Glu			Gly
	590	595			600
Asp Val Phe Pro	Gly Gln Val Lys Arg	Lys Tyr			
	605	610			

<210> 9

<211> 1469

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 60203310CD1

<400> 9

Met Trp Pro Ser Gln	Leu Leu Ile Phe	Met Met Leu Leu Ala	Pro
1	5	10	15
Ile Ile His Ala Phe	Ser Arg Ala Pro	Ile Pro Met Ala Val	Val
	20	25	30
Arg Arg Glu Leu Ser	Cys Glu Ser Tyr	Pro Ile Glu Leu Arg	Cys
	35	40	45
Pro Gly Thr Asp Val	Ile Met Ile Glu	Ser Ala Asn Tyr Gly	Arg
	50	55	60
Thr Asp Asp Lys Ile	Cys Asp Ser Asp	Pro Ala Gln Met Glu	Asn
	65	70	75
Ile Arg Cys Tyr Leu	Pro Asp Ala Tyr	Lys Ile Met Ser Gln	Arg
	80	85	90
Cys Asn Asn Arg Thr	Gln Cys Ala Val	Val Ala Gly Pro Asp	Val
	95	100	105
Phe Pro Asp Pro Cys	Pro Gly Thr Tyr	Lys Tyr Leu Glu Val	Gln
	110	115	120
Tyr Glu Cys Val Pro	Tyr Lys Val Glu	Gln Lys Val Phe Leu	Cys
	125	130	135
Pro Gly Leu Leu Lys	Gly Val Tyr Gln	Ser Glu His Leu Phe	Glu
	140	145	150
Ser Asp His Gln Ser	Gly Ala Trp Cys	Lys Asp Pro Leu Gln	Ala
	155	160	165
Ser Asp Lys Ile Tyr	Tyr Met Pro Trp	Thr Pro Tyr Arg Thr	Asp
	170	175	180
Thr Leu Thr Glu Tyr	Ser Ser Lys Asp	Asp Phe Ile Ala Gly	Arg
	185	190	195
Pro Thr Thr Thr Tyr	Lys Leu Pro His	Arg Val Asp Gly Thr	Gly
	200	205	210
Phe Val Val Tyr Asp	Gly Ala Leu Phe	Phe Asn Lys Glu Arg	Thr
	215	220	225
Arg Asn Ile Val Lys	Phe Asp Leu Arg	Thr Arg Ile Lys Ser	Gly
	230	235	240
Glu Ala Ile Ile Ala	Asn Ala Asn Tyr	His Asp Thr Ser Pro	Tyr
	245	250	255
Arg Trp Gly Gly Lys	Ser Asp Ile Asp	Leu Ala Val Asp Glu	Asn
	260	265	270
Gly Leu Trp Val Ile	Tyr Ala Thr Glu	Gln Asn Asn Gly Lys	Ile

	275	280	285
Val Ile Ser Gln	Leu Asn Pro Tyr Thr	Leu Arg Ile Glu Gly Thr	
	290	295	300
Trp Asp Thr Ala	Tyr Asp Lys Arg Ser	Ala Ser Asn Ala Phe Met	
	305	310	315
Ile Cys Gly Ile	Leu Tyr Val Val Lys	Ser Val Tyr Glu Asp Asp	
	320	325	330
Asp Asn Glu Ala	Thr Gly Asn Lys Ile	Asp Tyr Ile Tyr Asn Thr	
	335	340	345
Asp Gln Ser Lys	Asp Ser Leu Val Asp	Val Pro Phe Pro Asn Ser	
	350	355	360
Tyr Gln Tyr Ile	Ala Ala Val Asp Tyr	Asn Pro Arg Asp Asn Leu	
	365	370	375
Leu Tyr Val Trp	Asn Asn Tyr His Val	Val Lys Tyr Ser Leu Asp	
	380	385	390
Phe Gly Pro Leu	Asp Ser Arg Ser Gly	Gln Ala His His Gly Gln	
	395	400	405
Val Ser Tyr Ile	Ser Pro Pro Ile His	Leu Asp Ser Glu Leu Glu	
	410	415	420
Arg Pro Ser Val	Lys Asp Ile Ser Thr	Thr Gly Pro Leu Gly Met	
	425	430	435
Gly Ser Thr Thr	Thr Ser Thr Thr Leu	Arg Thr Thr Thr Leu Ser	
	440	445	450
Pro Gly Arg Ser	Thr Thr Pro Ser Val	Ser Gly Arg Arg Asn Arg	
	455	460	465
Ser Thr Ser Thr	Pro Ser Pro Ala Val	Glu Val Leu Asp Asp Met	
	470	475	480
Thr Thr His Leu	Pro Ser Ala Ser Ser	Gln Ile Pro Ala Leu Glu	
	485	490	495
Glu Ser Cys Glu	Ala Val Glu Ala Arg	Glu Ile Met Trp Phe Lys	
	500	505	510
Thr Arg Gln Gly	Gln Ile Ala Lys Gln	Pro Cys Pro Ala Gly Thr	
	515	520	525
Ile Gly Val Ser	Thr Tyr Leu Cys Leu	Ala Pro Asp Gly Ile Trp	
	530	535	540
Asp Pro Gln Gly	Pro Asp Leu Ser Asn	Cys Ser Ser Pro Trp Val	
	545	550	555
Asn His Ile Thr	Gln Lys Leu Lys Ser	Gly Glu Thr Ala Ala Asn	
	560	565	570
Ile Ala Arg Glu	Leu Ala Glu Gln Thr	Arg Asn His Leu Asn Ala	
	575	580	585
Gly Asp Ile Thr	Tyr Ser Val Arg Ala	Met Asp Gln Leu Val Gly	
	590	595	600
Leu Leu Asp Val	Gln Leu Arg Asn Leu	Thr Pro Gly Gly Lys Asp	
	605	610	615
Ser Ala Ala Arg	Ser Leu Asn Lys Leu	Gln Lys Arg Glu Arg Ser	
	620	625	630
Cys Arg Ala Tyr	Val Gln Ala Met Val	Glu Thr Val Asn Asn Leu	
	635	640	645
Leu Gln Pro Gln	Ala Leu Asn Ala Trp	Arg Asp Leu Thr Thr Ser	
	650	655	660
Asp Gln Leu Arg	Ala Ala Thr Met Leu	Leu His Thr Val Glu Glu	
	665	670	675
Ser Ala Phe Val	Leu Ala Asp Asn Leu	Leu Lys Thr Asp Ile Val	
	680	685	690
Arg Glu Asn Thr	Asp Asn Ile Lys Leu	Glu Val Ala Arg Leu Ser	
	695	700	705
Thr Glu Gly Asn	Leu Glu Asp Leu Lys	Phe Pro Glu Asn Met Gly	
	710	715	720
His Gly Ser Thr	Ile Gln Leu Ser Ala	Asn Thr Leu Lys Gln Asn	
	725	730	735
Gly Arg Asn Gly	Glu Ile Arg Val Ala	Phe Val Leu Tyr Asn Asn	
	740	745	750

Leu Gly Pro Tyr	Leu Ser Thr Glu Asn Ala	Ser Met Lys Leu Gly	
755	760	765	
Thr Glu Ala Leu	Ser Thr Asn His Ser Val	Ile Val Asn Ser Pro	
770	775	780	
Val Ile Thr Ala	Ala Ile Asn Lys Glu Phe	Ser Asn Lys Val Tyr	
785	790	795	
Leu Ala Asp Pro	Val Val Phe Thr Val Lys	His Ile Lys Gln Ser	
800	805	810	
Glu Glu Asn Phe	Asn Pro Asn Cys Ser Phe	Trp Ser Tyr Ser Lys	
815	820	825	
Arg Thr Met Thr	Gly Tyr Trp Ser Thr Gln	Gly Cys Arg Leu Leu	
830	835	840	
Thr Thr Asn Lys	Thr His Thr Thr Cys Ser	Cys Asn His Leu Thr	
845	850	855	
Asn Phe Ala Val	Leu Met Ala His Val Glu	Val Lys His Ser Asp	
860	865	870	
Ala Val His Asp	Leu Leu Leu Asp Val Ile	Thr Trp Val Gly Ile	
875	880	885	
Leu Leu Ser Leu	Val Cys Leu Leu Ile Cys	Ile Phe Thr Phe Cys	
890	895	900	
Phe Phe Arg Gly	Leu Gln Ser Asp Arg Asn	Thr Ile His Lys Asn	
905	910	915	
Leu Cys Ile Ser	Leu Phe Val Ala Glu Leu	Leu Phe Leu Ile Gly	
920	925	930	
Ile Asn Arg Thr	Asp Gln Pro Ile Ala Cys	Ala Val Phe Ala Ala	
935	940	945	
Leu Leu His Phe	Phe Phe Leu Ala Ala Phe	Thr Trp Met Phe Leu	
950	955	960	
Glu Gly Val Gln	Leu Tyr Ile Met Leu Val	Glu Val Phe Glu Ser	
965	970	975	
Glu His Ser Arg	Arg Lys Tyr Phe Tyr Leu	Val Gly Tyr Gly Met	
980	985	990	
Pro Ala Leu Ile	Val Ala Val Ser Ala Ala	Val Asp Tyr Arg Ser	
995	1000	1005	
Tyr Gly Thr Asp	Lys Val Cys Trp Leu Arg	Leu Asp Thr Tyr Phe	
1010	1015	1020	
Ile Trp Ser Phe	Ile Gly Pro Ala Thr Leu	Ile Ile Met Leu Asn	
1025	1030	1035	
Val Ile Phe Leu	Gly Ile Ala Leu Tyr Lys	Met Val His His Thr	
1040	1045	1050	
Ala Ile Leu Lys	Pro Glu Ser Gly Cys Leu	Asp Asn Ile Asn Tyr	
1055	1060	1065	
Glu Asp Asn Arg	Pro Phe Ile Lys Ser Trp	Val Ile Gly Ala Ile	
1070	1075	1080	
Ala Leu Leu Cys	Leu Leu Gly Leu Thr Trp	Ala Phe Gly Leu Met	
1085	1090	1095	
Tyr Ile Asn Glu	Ser Thr Val Ile Met Ala	Tyr Leu Phe Thr Ile	
1100	1105	1110	
Phe Asn Ser Leu	Gln Gly Met Phe Ile Phe	Ile Phe His Cys Val	
1115	1120	1125	
Leu Gln Lys Lys	Val Arg Lys Glu Tyr Gly	Lys Cys Leu Arg Thr	
1130	1135	1140	
His Cys Cys Ser	Gly Lys Ser Thr Glu Ser	Ser Ile Gly Ser Gly	
1145	1150	1155	
Lys Thr Ser Gly	Ser Arg Thr Pro Gly Arg	Tyr Ser Thr Gly Ser	
1160	1165	1170	
Gln Ser Arg Ile	Arg Arg Met Trp Asn Asp	Thr Val Arg Lys Gln	
1175	1180	1185	
Ser Glu Ser Ser	Phe Ile Thr Gly Asp Ile	Asn Ser Ser Ala Ser	
1190	1195	1200	
Leu Asn Arg Glu	Gly Leu Leu Asn Asn Ala	Arg Asp Thr Ser Val	
1205	1210	1215	
Met Asp Thr Leu	Pro Leu Asn Gly Asn His	Gly Asn Ser Tyr Ser	

	1220		1225		1230
Ile Ala Ser Gly Glu Tyr Leu Ser Asn Cys Val Gln Ile Ile Asp					
	1235		1240		1245
Arg Gly Tyr Asn His Asn Glu Thr Ala Leu Glu Lys Lys Ile Leu					
	1250		1255		1260
Lys Glu Leu Thr Ser Asn Tyr Ile Pro Ser Tyr Leu Asn Asn His					
	1265		1270		1275
Glu Arg Ser Ser Glu Gln Asn Arg Asn Leu Met Asn Lys Leu Val					
	1280		1285		1290
Asn Asn Leu Gly Ser Gly Arg Glu Asp Asp Ala Ile Val Leu Asp					
	1295		1300		1305
Asp Ala Thr Ser Phe Asn His Glu Glu Ser Leu Gly Leu Glu Leu					
	1310		1315		1320
Ile His Glu Glu Ser Asp Ala Pro Leu Leu Pro Pro Arg Val Tyr					
	1325		1330		1335
Ser Thr Glu Asn His Gln Pro His His Tyr Thr Arg Arg Arg Ile					
	1340		1345		1350
Pro Gln Asp His Ser Glu Ser Phe Phe Pro Leu Leu Thr Asn Glu					
	1355		1360		1365
His Thr Glu Asp Leu Gln Ser Pro His Arg Asp Ser Leu Tyr Thr					
	1370		1375		1380
Ser Met Pro Thr Leu Ala Gly Val Ala Ala Thr Glu Ser Val Thr					
	1385		1390		1395
Thr Ser Thr Gln Thr Glu Pro Pro Pro Ala Lys Cys Gly Asp Ala					
	1400		1405		1410
Glu Asp Val Tyr Tyr Lys Ser Met Pro Asn Leu Gly Ser Arg Asn					
	1415		1420		1425
His Val His Gln Leu His Thr Tyr Tyr Gln Leu Gly Arg Gly Ser					
	1430		1435		1440
Ser Asp Gly Phe Ile Val Pro Pro Asn Lys Asp Gly Thr Pro Pro					
	1445		1450		1455
Glu Gly Ser Ser Lys Gly Pro Ala His Leu Val Thr Ser Leu					
	1460		1465		

<210> 10

<211> 469

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7477349CD1

<400> 10

Met Asp Pro Ser Val Val Ser Asn Glu Tyr Tyr Asp Val Ala His					
1	5		10		15
Gly Ala Lys Asp Pro Val Val Pro Thr Ser Leu Gln Asp Ile Thr					
	20		25		30
Ala Val Leu Gly Thr Glu Ala Tyr Thr Glu Glu Asp Lys Ser Met					
	35		40		45
Val Ser His Ala Gln Lys Ser Gln His Ser Cys Leu Ser His Ser					
	50		55		60
Arg Trp Leu Arg Ser Pro Gln Val Thr Gly Gly Ser Trp Asp Leu					
	65		70		75
Arg Ile Arg Pro Ser Lys Asp Ser Ser Ser Phe Arg Gln Ala Gln					
	80		85		90
Cys Leu Arg Lys Asp Pro Gly Ala Asn Asn His Leu Glu Ser Gln					
	95		100		105
Gly Val Arg Gly Thr Ala Gly Asp Ala Asp Arg Glu Leu Arg Gly					
	110		115		120
Pro Ser Glu Lys Ala Thr Ala Gly Gln Pro Arg Val Thr Leu Leu					
	125		130		135
Pro Thr Pro Asn Val Ser Gly Leu Ser Gln Glu Phe Glu Ser His					

Trp	Pro	Glu	Ile	Ala	Glu	Arg	Ser	Pro	Cys	Val	Ala	Gly	Val	Ile
				140					145					150
Pro	Val	Ile	Tyr	Tyr	Ser	Val	Leu	Leu	Gly	Leu	Gly	Leu	Pro	Val
				155					160					165
Ser	Leu	Leu	Thr	Ala	Val	Ala	Leu	Ala	Arg	Leu	Ala	Thr	Arg	Thr
				170					175					180
Arg	Arg	Pro	Ser	Tyr	Tyr	Tyr	Leu	Leu	Ala	Leu	Thr	Ala	Ser	Asp
				185					190					195
Ile	Ile	Ile	Gln	Val	Val	Ile	Val	Phe	Ala	Gly	Phe	Leu	Leu	Gln
				200					205					210
Gly	Ala	Val	Leu	Ala	Arg	Gln	Val	Pro	Gln	Ala	Val	Val	Arg	Thr
				215					220					225
Ala	Asn	Ile	Leu	Glu	Phe	Ala	Ala	Asn	His	Ala	Ser	Val	Trp	Ile
				230					235					240
Ala	Ile	Leu	Leu	Thr	Val	Asp	Arg	Tyr	Thr	Ala	Leu	Cys	His	Pro
				245					250					255
Leu	His	His	Arg	Ala	Ala	Ser	Ser	Pro	Gly	Arg	Thr	Arg	Arg	Ala
				260					265					270
Ile	Ala	Ala	Val	Leu	Ser	Ala	Ala	Leu	Leu	Thr	Gly	Ile	Pro	Phe
				275					280					285
Tyr	Trp	Trp	Leu	Asp	Met	Trp	Arg	Asp	Thr	Asp	Ser	Pro	Arg	Thr
				290					295					300
Leu	Asp	Glu	Val	Leu	Lys	Trp	Ala	His	Cys	Leu	Thr	Val	Tyr	Phe
				305					310					315
Ile	Pro	Cys	Gly	Val	Phe	Leu	Val	Thr	Asn	Ser	Ala	Ile	Ile	His
				320					325					330
Arg	Leu	Arg	Arg	Arg	Gly	Arg	Ser	Gly	Leu	Gln	Pro	Arg	Val	Gly
				335					340					345
Lys	Ser	Thr	Ala	Ile	Leu	Leu	Gly	Ile	Thr	Thr	Leu	Phe	Thr	Leu
				350					355					360
Leu	Trp	Ala	Pro	Arg	Val	Phe	Val	Met	Leu	Tyr	His	Met	Tyr	Val
				365					370					375
Ala	Pro	Val	His	Arg	Asp	Trp	Arg	Val	His	Leu	Ala	Leu	Asp	Val
				380					385					390
Ala	Asn	Met	Val	Ala	Met	Leu	His	Thr	Ala	Ala	Asn	Phe	Gly	Leu
				395					400					405
Tyr	Cys	Phe	Val	Ser	Lys	Thr	Phe	Arg	Ala	Thr	Val	Arg	Gln	Val
				410					415					420
Ile	His	Asp	Ala	Tyr	Leu	Pro	Cys	Thr	Leu	Ala	Ser	Gln	Pro	Glu
				425					430					435
Gly	Met	Ala	Ala	Lys	Pro	Val	Met	Glu	Pro	Pro	Gly	Leu	Pro	Thr
				440					445					450
Gly	Ala	Glu	Val						455					460
									460					465

<210> 11
 <211> 335
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 55002225CD1

<400> 11
 Met Asn Pro Phe His Ala Ser Cys Trp Asn Thr Ser Ala Glu Leu
 1 5 10 15
 Leu Asn Lys Ser Trp Asn Lys Glu Phe Ala Tyr Gln Thr Ala Ser
 20 25 30
 Val Val Asp Thr Val Ile Leu Pro Ser Met Ile Gly Ile Ile Cys
 35 40 45
 Ser Thr Gly Leu Val Gly Asn Ile Leu Ile Val Phe Thr Ile Ile

Arg	Ser	Arg	Lys	Lys	Thr	Val	Pro	Asp	Ile	Tyr	Ile	Cys	Asn	Leu	
				65					55						60
Ala	Val	Ala	Asp	Leu	Val	His	Ile	Val	Gly	Met	Pro	Phe	Leu	Ile	
				80					70						75
His	Gln	Trp	Ala	Arg	Gly	Gly	Glu	Trp	Val	Phe	Gly	Gly	Pro	Leu	
				95					85						90
Cys	Thr	Ile	Ile	Thr	Ser	Leu	Asp	Thr	Cys	Asn	Gln	Phe	Ala	Cys	
				110					100						105
Ser	Ala	Ile	Met	Thr	Val	Met	Ser	Val	Asp	Arg	Tyr	Phe	Ala	Leu	
				125					115						120
Val	Gln	Pro	Phe	Arg	Leu	Thr	Arg	Trp	Arg	Thr	Arg	Tyr	Lys	Thr	
				140					130						135
Ile	Arg	Ile	Asn	Leu	Gly	Leu	Trp	Ala	Ala	Ser	Phe	Ile	Leu	Ala	
				155					145						150
Leu	Pro	Val	Trp	Val	Tyr	Ser	Lys	Val	Ile	Lys	Phe	Lys	Asp	Gly	
				170					160						165
Val	Glu	Ser	Cys	Ala	Phe	Asp	Leu	Thr	Ser	Pro	Asp	Asp	Val	Leu	
				185					175						180
Trp	Tyr	Thr	Leu	Tyr	Leu	Thr	Ile	Thr	Thr	Phe	Phe	Phe	Pro	Leu	
				200					190						195
Pro	Leu	Ile	Leu	Val	Cys	Tyr	Ile	Leu	Ile	Leu	Cys	Tyr	Thr	Trp	
				215					205						210
Glu	Met	Tyr	Gln	Gln	Asn	Lys	Asp	Ala	Arg	Cys	Cys	Asn	Pro	Ser	
				230					220						225
Val	Pro	Lys	Gln	Arg	Val	Met	Lys	Leu	Thr	Lys	Met	Val	Leu	Val	
				245					235						240
Leu	Val	Val	Val	Phe	Ile	Leu	Ser	Ala	Ala	Pro	Tyr	His	Val	Ile	
				260					250						255
Gln	Leu	Val	Asn	Leu	Gln	Met	Glu	Gln	Pro	Thr	Leu	Ala	Phe	Tyr	
				275					265						270
Val	Gly	Tyr	Tyr	Leu	Ser	Ile	Cys	Leu	Ser	Tyr	Ala	Ser	Ser	Ser	
				290					280						285
Ile	Asn	Pro	Phe	Leu	Tyr	Ile	Leu	Leu	Ser	Gly	Thr	Pro	Gln	Ile	
				305					295						300
Gln	Arg	Arg	Ala	Thr	Glu	Lys	Glu	Ile	Asn	Asn	Met	Gly	Asn	Thr	
				320					310						315
Leu	Lys	Ser	His	Phe					325						330
				335											

<210> 12

<211> 630

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7475686CD1

<400> 12

Met	Arg	Leu	Gly	Pro	Val	Pro	Ala	Arg	Ala	Arg	Ala	Leu	Leu	Ser	
1				5					10					15	
Trp	Val	Arg	Gly	Leu	Glu	Ser	Arg	Gly	Gly	Glu	Trp	Thr	Lys	Cys	
				20					25					30	
Ile	Val	Gln	Leu	Gly	His	Leu	Leu	Ala	Thr	Gln	His	Pro	Ala	Ala	
				35					40					45	
Pro	Thr	Cys	Gly	Val	Val	Ser	Ser	Ala	Leu	Val	Met	His	Ser	Thr	
				50					55					60	
Asp	Val	Cys	Leu	Ala	Pro	Thr	Met	His	Gln	Ala	Leu	Asp	Trp	Ala	
				65					70					75	
Ala	Gly	Ile	Trp	Phe	Thr	Gly	Arg	Leu	Gly	Leu	Arg	Glu	His	Lys	
				80					85					90	
Ser	Leu	Ala	Gln	Gly	Asp	Ser	Val	Cys	Pro	Cys	Glu	Ser	Glu	Leu	

	95		100		105
Gly Asp Phe Gln Val Tyr Gly Leu Val Ser Thr Glu Gly Val Val					
	110		115		120
Ser Cys Phe Gly Glu Lys Thr Pro Gln His Pro Gly Pro Pro Ala					
	125		130		135
Ser Leu Ser Leu Ala Asn Arg Cys His Asn Val Val Thr Ala Val					
	140		145		150
Gly Ala Trp Pro Ala His Gly Ser Ile Leu Gly Asn Val Pro Glu					
	155		160		165
Ala Pro Val Gly Ala Asp Val Leu Gly Ala Gly Gly Cys Asp Trp					
	170		175		180
Ala Asp Lys Glu Ala Leu Ala Pro Gly Gln Arg Ala Lys Val His					
	185		190		195
Ile Leu Leu Glu Ser Ser Gly Gln Ser Asp Pro Ser Tyr Ala Val					
	200		205		210
Leu Pro Asp Ser Trp Ala Ala Thr Glu Gly Phe Pro Thr Tyr Arg					
	215		220		225
Ser Gln Val Ser Ser Pro Arg Ile Pro Gly Ser Ser Ile Trp Leu					
	230		235		240
Gly Ser Gly Ser Gly Trp Pro Ile Leu Gly Glu Leu Arg Glu Cys					
	245		250		255
Asp Gln Met Phe Ser Cys Met Leu Pro Thr Gly Cys Ala Ser Phe					
	260		265		270
Gln Asp Pro Gly Arg Tyr Gly Asp Tyr Asp Leu Pro Met Asp Glu					
	275		280		285
Asp Glu Asp Met Thr Lys Thr Arg Thr Phe Phe Ala Ala Lys Ile					
	290		295		300
Val Ile Gly Ile Ala Leu Ala Gly Ile Met Leu Val Cys Gly Ile					
	305		310		315
Gly Asn Phe Val Phe Ile Ala Ala Leu Thr Arg Tyr Lys Lys Leu					
	320		325		330
Arg Asn Leu Thr Asn Leu Leu Ile Ala Asn Leu Ala Ile Ser Asp					
	335		340		345
Phe Leu Val Ala Ile Ile Cys Cys Pro Phe Glu Met Asp Tyr Tyr					
	350		355		360
Val Val Arg Gln Leu Ser Trp Glu His Gly His Val Leu Cys Ala					
	365		370		375
Ser Val Asn Tyr Leu Arg Thr Val Ser Leu Tyr Val Ser Thr Asn					
	380		385		390
Ala Leu Leu Ala Ile Ala Ile Asp Arg Tyr Leu Ala Ile Val His					
	395		400		405
Pro Leu Lys Pro Arg Met Asn Tyr Gln Thr Ala Ser Phe Leu Ile					
	410		415		420
Ala Leu Val Trp Met Val Ser Ile Leu Ile Ala Ile Pro Ser Ala					
	425		430		435
Tyr Phe Ala Thr Glu Thr Val Leu Phe Ile Val Lys Ser Gln Glu					
	440		445		450
Lys Ile Phe Cys Gly Gln Ile Trp Pro Val Asp Gln Gln Leu Tyr					
	455		460		465
Tyr Lys Ser Tyr Phe Leu Phe Ile Phe Gly Val Glu Phe Val Gly					
	470		475		480
Pro Val Val Thr Met Thr Leu Cys Tyr Ala Arg Ile Ser Arg Glu					
	485		490		495
Leu Trp Phe Lys Ala Val Pro Gly Phe Gln Thr Glu Gln Ile Arg					
	500		505		510
Lys Arg Leu Arg Cys Arg Arg Lys Thr Val Leu Val Leu Met Cys					
	515		520		525
Ile Leu Thr Ala Tyr Val Leu Cys Trp Ala Pro Phe Tyr Gly Phe					
	530		535		540
Thr Ile Val Arg Asp Phe Phe Pro Thr Val Phe Val Lys Glu Lys					
	545		550		555
His Tyr Leu Thr Ala Phe Tyr Val Val Glu Cys Ile Ala Met Ser					
	560		565		570

Asn	Ser	Met	Ile	Asn	Thr	Val	Cys	Phe	Val	Thr	Val	Lys	Asn	Asn	
				575					580					585	
Thr	Met	Lys	Tyr	Phe	Lys	Lys	Met	Met	Leu	Leu	His	Trp	Arg	Pro	
				590					595					600	
Ser	Gln	Arg	Gly	Ser	Lys	Ser	Ser	Ala	Asp	Leu	Asp	Leu	Arg	Thr	
				605					610					615	
Asn	Gly	Val	Pro	Thr	Thr	Glu	Glu	Val	Asp	Cys	Ile	Arg	Leu	Lys	
				620					625					630	

<210> 13

<211> 695

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7482007CD1

<400> 13

Met	Lys	Met	Lys	Ser	Gln	Ala	Thr	Met	Ile	Cys	Cys	Leu	Val	Phe	
1				5					10					15	
Phe	Leu	Ser	Thr	Glu	Cys	Ser	His	Tyr	Arg	Ser	Lys	Ile	His	Leu	
				20					25					30	
Lys	Ala	Gly	Asp	Lys	Leu	Gln	Ser	Pro	Glu	Gly	Lys	Pro	Lys	Thr	
				35					40					45	
Gly	Arg	Ile	Gln	Glu	Lys	Cys	Glu	Gly	Pro	Cys	Ile	Ser	Ser	Ser	
				50					55					60	
Asn	Cys	Ser	Gln	Pro	Cys	Ala	Lys	Asp	Phe	His	Gly	Glu	Ile	Gly	
				65					70					75	
Phe	Thr	Cys	Asn	Gln	Lys	Lys	Trp	Gln	Lys	Ser	Ala	Glu	Thr	Cys	
				80					85					90	
Thr	Ser	Leu	Ser	Val	Glu	Lys	Leu	Phe	Lys	Asp	Ser	Thr	Gly	Ala	
				95					100					105	
Ser	Arg	Leu	Ser	Val	Ala	Ala	Pro	Ser	Ile	Pro	Leu	His	Ile	Leu	
				110					115					120	
Asp	Phe	Arg	Ala	Pro	Glu	Thr	Ile	Glu	Ser	Val	Ala	Gln	Gly	Ile	
				125					130					135	
Arg	Lys	Asn	Cys	Pro	Phe	Asp	Tyr	Ala	Cys	Ile	Thr	Asp	Met	Val	
				140					145					150	
Lys	Ser	Ser	Glu	Thr	Thr	Ser	Gly	Asn	Ile	Ala	Phe	Ile	Val	Glu	
				155					160					165	
Leu	Leu	Lys	Asn	Ile	Ser	Thr	Asp	Leu	Ser	Asp	Asn	Val	Thr	Arg	
				170					175					180	
Glu	Lys	Met	Lys	Ser	Tyr	Ser	Glu	Val	Ala	Asn	His	Ile	Leu	Asp	
				185					190					195	
Thr	Ala	Ala	Ile	Ser	Asn	Trp	Ala	Phe	Ile	Pro	Asn	Lys	Asn	Ala	
				200					205					210	
Ser	Ser	Asp	Leu	Leu	Gln	Ser	Val	Asn	Leu	Phe	Ala	Arg	Gln	Leu	
				215					220					225	
His	Ile	His	Asn	Asn	Ser	Glu	Asn	Ile	Val	Asn	Glu	Leu	Phe	Ile	
				230					235					240	
Gln	Thr	Lys	Gly	Phe	His	Ile	Asn	His	Asn	Thr	Ser	Glu	Lys	Ser	
				245					250					255	
Leu	Asn	Phe	Ser	Met	Ser	Met	Asn	Asn	Thr	Thr	Glu	Asp	Ile	Leu	
				260					265					270	
Gly	Met	Val	Gln	Ile	Pro	Arg	Gln	Glu	Leu	Arg	Lys	Leu	Trp	Pro	
				275					280					285	
Asn	Ala	Ser	Gln	Ala	Ile	Ser	Ile	Ala	Phe	Pro	Thr	Leu	Gly	Ala	
				290					295					300	
Ile	Leu	Arg	Glu	Ala	His	Leu	Gln	Asn	Val	Ser	Leu	Pro	Arg	Gln	
				305					310					315	
Val	Asn	Gly	Leu	Val	Leu	Ser	Val	Val	Leu	Pro	Glu	Arg	Leu	Gln	

Glu Ile Ile Leu	320	325	330
Thr Phe Glu Lys Ile		Asn Lys Thr Arg Asn	Ala
335	340		345
Arg Ala Gln Cys	Val Gly Trp His Ser	Lys Lys Arg Arg Trp	Asp
350	355		360
Glu Lys Ala Cys	Gln Met Met Leu Asp	Ile Arg Asn Glu Val	Lys
365	370		375
Cys Arg Cys Asn	Tyr Thr Ser Val Val	Met Ser Phe Ser Ile	Leu
380	385		390
Met Ser Ser Lys	Ser Met Thr Asp Lys	Val Leu Asp Tyr Ile	Thr
395	400		405
Cys Ile Gly Leu	Ser Val Ser Ile Leu	Ser Leu Val Leu Cys	Leu
410	415		420
Ile Ile Glu Ala	Thr Val Trp Ser Arg	Val Val Val Thr Glu	Ile
425	430		435
Ser Tyr Met Arg	His Val Cys Ile Val	Asn Ile Ala Val Ser	Leu
440	445		450
Leu Thr Ala Asn	Val Trp Phe Ile Ile	Gly Ser His Phe Asn	Ile
455	460		465
Lys Ala Gln Asp	Tyr Asn Met Cys Val	Ala Val Thr Phe Phe	Ser
470	475		480
His Phe Phe Tyr	Leu Ser Leu Phe Phe	Trp Ile Leu Phe Lys	Ala
485	490		495
Leu Leu Ile Ile	Tyr Gly Ile Leu Val	Ile Phe Arg Arg Met	Met
500	505		510
Lys Ser Arg Met	Met Val Ile Gly Phe	Ala Ile Gly Tyr Gly	Cys
515	520		525
Pro Leu Ile Ile	Ala Val Thr Thr Val	Ala Ile Thr Gly Pro	Val
530	535		540
Lys Gly Tyr Met	Arg Pro Glu Ala Cys	Trp Leu Asn Trp Asp	Asn
545	550		555
Thr Lys Ala Leu	Leu Ala Phe Ala Ile	Pro Ala Phe Val Ile	Val
560	565		570
Ala Val Asn Leu	Ile Val Val Leu Val	Val Ala Val Asn Thr	Gln
575	580		585
Arg Pro Ser Ile	Gly Ser Ser Lys Ser	Gln Asp Val Val Ile	Ile
590	595		600
Met Arg Ile Ser	Lys Asn Val Ala Ile	Leu Thr Pro Leu Leu	Gly
605	610		615
Leu Thr Trp Gly	Phe Gly Ile Ala Thr	Leu Ile Glu Gly Thr	Ser
620	625		630
Leu Thr Phe His	Ile Ile Phe Ala Leu	Leu Asn Ala Phe Gln	Gly
635	640		645
Phe Phe Ile Leu	Leu Phe Gly Thr Ile	Met Asp His Lys Ile	Arg
650	655		660
Asp Ala Leu Arg	Met Arg Met Ser Ser	Leu Lys Gly Lys Ser	Arg
665	670		675
Ala Ala Glu Asn	Ala Ser Leu Gly Pro	Thr Asn Gly Ser Lys	Leu
680	685		690
Met Asn Arg Gln	Gly		
695			

<210> 14

<211> 633

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 6769042CD1

<400> 14

Met Tyr Phe Thr Ala Ala Ile Gly Lys His Ala Leu Leu Ser Ser

1	5	10	15
Thr Leu Pro Ser	Leu Phe Met Thr Ser	Thr Ala Ser Pro Val	Met
	20	25	30
Pro Thr Asp Ala	Tyr His Pro Ile Ile	Thr Asn Leu Thr Glu	Glu
	35	40	45
Arg Lys Thr Phe	Gln Ser Pro Gly Val	Ile Leu Ser Tyr Leu	Gln
	50	55	60
Asn Val Ser Leu	Ser Leu Pro Ser Lys	Ser Leu Ser Glu Gln	Thr
	65	70	75
Ala Leu Asn Leu	Thr Lys Thr Phe Leu	Lys Ala Val Gly Glu	Ile
	80	85	90
Leu Leu Leu Pro	Gly Trp Ile Ala Leu	Ser Glu Asp Ser Ala	Val
	95	100	105
Val Leu Ser Leu	Ile Asp Thr Ile Asp	Thr Val Met Gly His	Val
	110	115	120
Ser Ser Asn Leu	His Gly Ser Thr Pro	Gln Val Thr Val Glu	Gly
	125	130	135
Ser Ser Ala Met	Ala Glu Phe Ser Val	Ala Lys Ile Leu Pro	Lys
	140	145	150
Thr Val Asn Ser	Ser His Tyr Arg Phe	Pro Ala His Gly Gln	Ser
	155	160	165
Phe Ile Gln Ile	Pro His Glu Ala Phe	His Arg His Ala Trp	Ser
	170	175	180
Thr Val Val Gly	Leu Leu Tyr His Ser	Met His Tyr Tyr Leu	Asn
	185	190	195
Asn Ile Trp Pro	Ala His Thr Lys Ile	Ala Glu Ala Met His	His
	200	205	210
Gln Asp Cys Leu	Leu Phe Ala Thr Ser	His Leu Ile Ser Leu	Glu
	215	220	225
Val Ser Pro Pro	Pro Thr Leu Ser Gln	Asn Leu Ser Gly Ser	Pro
	230	235	240
Leu Ile Thr Val	His Leu Lys His Arg	Leu Thr Arg Lys Gln	His
	245	250	255
Ser Glu Ala Thr	Asn Ser Ser Asn Arg	Val Phe Val Tyr Cys	Ala
	260	265	270
Phe Leu Asp Phe	Ser Ser Gly Glu Gly	Val Trp Ser Asn His	Gly
	275	280	285
Cys Ala Leu Thr	Arg Gly Asn Leu Thr	Tyr Ser Val Cys Arg	Cys
	290	295	300
Thr His Leu Thr	Asn Phe Ala Ile Leu	Met Gln Val Val Pro	Leu
	305	310	315
Glu Leu Ala Arg	Gly His Gln Val Ala	Leu Ser Ser Ile Ser	Tyr
	320	325	330
Val Gly Cys Ser	Leu Ser Val Leu Cys	Leu Val Ala Thr Leu	Val
	335	340	345
Thr Phe Ala Val	Leu Ser Ser Val Ser	Thr Ile Arg Asn Gln	Arg
	350	355	360
Tyr His Ile His	Ala Asn Leu Ser Phe	Ala Val Leu Val Ala	Gln
	365	370	375
Val Leu Leu Leu	Ile Ser Phe Arg Leu	Glu Pro Gly Thr Thr	Pro
	380	385	390
Cys Gln Val Met	Ala Val Leu Leu His	Tyr Phe Phe Leu Ser	Ala
	395	400	405
Phe Ala Trp Met	Leu Val Glu Gly Leu	His Leu Tyr Ser Met	Val
	410	415	420
Ile Lys Val Phe	Gly Ser Glu Asp Ser	Lys His Arg Tyr Tyr	Tyr
	425	430	435
Gly Met Gly Trp	Gly Phe Pro Leu Leu	Ile Cys Ile Ile Ser	Leu
	440	445	450
Ser Phe Ala Met	Asp Ser Tyr Gly Thr	Ser Asn Asn Cys Trp	Leu
	455	460	465
Ser Leu Ala Ser	Gly Ala Ile Trp Ala	Phe Val Ala Pro Ala	Leu
	470	475	480

Phe	Val	Ile	Val	Val	Asn	Ile	Gly	Ile	Leu	Ile	Ala	Val	Thr	Arg
				485					490					495
Val	Ile	Ser	Gln	Ile	Ser	Ala	Asp	Asn	Tyr	Lys	Ile	His	Gly	Asp
				500					505					510
Pro	Ser	Ala	Phe	Lys	Leu	Thr	Ala	Lys	Ala	Val	Ala	Val	Leu	Leu
				515					520					525
Pro	Ile	Leu	Gly	Thr	Ser	Trp	Val	Phe	Gly	Val	Leu	Ala	Val	Asn
				530					535					540
Gly	Cys	Ala	Val	Val	Phe	Gln	Tyr	Met	Phe	Ala	Thr	Leu	Asn	Ser
				545					550					555
Leu	Gln	Gly	Leu	Phe	Ile	Phe	Leu	Phe	His	Cys	Leu	Leu	Asn	Ser
				560					565					570
Glu	Val	Arg	Ala	Ala	Phe	Lys	His	Lys	Ile	Lys	Val	Trp	Ser	Leu
				575					580					585
Thr	Ser	Ser	Ser	Ala	Arg	Thr	Ser	Asn	Ala	Lys	Pro	Phe	His	Ser
				590					595					600
Asp	Leu	Met	Asn	Gly	Thr	Arg	Pro	Gly	Met	Ala	Ser	Thr	Lys	Leu
				605					610					615
Ser	Pro	Trp	Asp	Lys	Ser	Ser	His	Ser	Ala	His	Arg	Val	Asp	Leu
				620					625					630
Ser	Ala	Val												

<210> 15

<211> 370

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7476053CD1

<400> 15

Met	Glu	Ala	Ala	Ser	Leu	Ser	Val	Ala	Thr	Ala	Gly	Val	Ala	Leu
1				5					10					15
Ala	Leu	Gly	Pro	Glu	Thr	Ser	Ser	Gly	Thr	Pro	Ser	Pro	Arg	Gly
				20					25					30
Ile	Leu	Gly	Ser	Thr	Pro	Ser	Gly	Ala	Val	Leu	Pro	Gly	Arg	Gly
				35					40					45
Pro	Pro	Phe	Ser	Val	Phe	Thr	Val	Leu	Val	Val	Thr	Leu	Leu	Val
				50					55					60
Leu	Leu	Ile	Ala	Ala	Thr	Phe	Leu	Trp	Asn	Leu	Leu	Val	Pro	Val
				65					70					75
Thr	Ile	Pro	Arg	Val	Arg	Ala	Phe	His	Arg	Val	Pro	His	Asn	Leu
				80					85					90
Val	Ala	Ser	Thr	Ala	Val	Ser	Asp	Glu	Leu	Val	Ala	Ala	Leu	Ala
				95					100					105
Met	Pro	Pro	Ser	Leu	Ala	Ser	Glu	Leu	Ser	Thr	Gly	Arg	Arg	Arg
				110					115					120
Leu	Leu	Gly	Arg	Ser	Leu	Cys	His	Val	Trp	Ile	Ser	Phe	Asp	Ala
				125					130					135
Leu	Cys	Cys	Pro	Ala	Gly	Leu	Gly	Asn	Val	Ala	Ala	Ile	Ala	Leu
				140					145					150
Gly	Arg	Asp	Gly	Ala	Ile	Thr	Arg	His	Leu	Gln	His	Thr	Leu	Arg
				155					160					165
Thr	Arg	Ser	Arg	Ala	Ser	Leu	Leu	Met	Ile	Ala	Leu	Ala	Arg	Val
				170					175					180
Pro	Ser	Ala	Leu	Ile	Ala	Leu	Ala	Pro	Leu	Leu	Phe	Gly	Arg	Gly
				185					190					195
Glu	Val	Cys	Asp	Ala	Arg	Leu	Gln	Arg	Cys	Gln	Val	Ser	Arg	Glu
				200					205					210
Pro	Ser	Tyr	Ala	Ala	Phe	Ser	Thr	Arg	Gly	Ala	Phe	His	Leu	Pro
				215					220					225

Leu Gly Val Val	Pro Phe Val Tyr Arg	Lys Ile Tyr Glu Ala	Ala
230		235	240
Lys Phe Arg Phe	Gly Arg Arg Arg Arg	Ala Val Leu Pro Leu	Pro
245		250	255
Ala Thr Met Gln	Val Lys Glu Ala Pro	Asp Glu Ala Glu Val	Val
260		265	270
Phe Thr Ala His	Cys Lys Ala Thr Val	Ser Phe Gln Val Ser	Gly
275		280	285
Asp Ser Trp Arg	Glu Gln Lys Glu Arg	Arg Ala Ala Met Met	Val
290		295	300
Gly Ile Leu Ile	Gly Val Phe Val Leu	Cys Trp Ile Pro Phe	Phe
305		310	315
Leu Thr Glu Leu	Ile Ser Pro Leu Cys	Ala Cys Ser Leu Pro	Pro
320		325	330
Ile Trp Lys Ser	Ile Phe Leu Trp Leu	Gly Tyr Ser Asn Ser	Phe
335		340	345
Phe Asn Pro Leu	Ile Tyr Thr Ala Phe	Asn Lys Asn Tyr Asn	Asn
350		355	360
Ala Phe Lys Ser	Leu Phe Thr Lys Gln	Arg	
365		370	

<210> 16

<211> 324

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7480410CD1

<400> 16

Met Gly Met Glu Gly	Leu Leu Gln Asn Ser	Thr Asn Phe Val Leu
1	5	10
Thr Gly Leu Ile Thr	His Pro Ala Phe Pro	Gly Leu Leu Phe Ala
20		25
Ile Val Phe Ser Ile	Phe Val Val Ala Ile	Thr Ala Asn Leu Val
35		40
Met Ile Leu Leu Ile	His Met Asp Ser Arg	Leu His Thr Pro Met
50		55
Tyr Phe Leu Leu Ser	Gln Leu Ser Ile Met	Asp Thr Ile Tyr Ile
65		70
Cys Ile Thr Val Pro	Lys Met Leu Gln Asp	Leu Leu Ser Lys Asp
80		85
Lys Thr Ile Ser Phe	Leu Gly Cys Ala Val	Gln Ile Phe Leu Tyr
95		100
Leu Thr Leu Ile Gly	Gly Glu Phe Phe Leu	Leu Gly Leu Met Ala
110		115
Tyr Asp Arg Tyr Val	Ala Val Cys Asn Pro	Leu Arg Tyr Pro Leu
125		130
Leu Met Asn Arg Arg	Val Cys Leu Phe Met	Val Val Gly Ser Trp
140		145
Val Gly Gly Ser Leu	Asp Gly Phe Met Leu	Thr Pro Val Thr Met
155		160
Ser Phe Pro Phe Cys	Arg Ser Arg Glu Ile	Asn His Phe Phe Cys
170		175
Glu Ile Pro Ala Val	Leu Lys Leu Ser Cys	Thr Asp Thr Ser Leu
185		190
Tyr Glu Thr Leu Met	Tyr Ala Cys Cys Val	Leu Met Leu Leu Ile
200		205
Pro Leu Ser Val Ile	Ser Val Ser Tyr Thr	His Ile Leu Leu Thr
215		220
Val His Arg Met Asn	Ser Ala Glu Gly Arg	Arg Lys Ala Phe Ala
230		235

```

Thr Cys Ser Ser His Ile Met Val Val Ser Val Phe Tyr Gly Ala
245 250 255
Ala Phe Tyr Thr Asn Val Leu Pro His Ser Tyr His Thr Pro Glu
260 265 270
Lys Asp Lys Val Val Ser Ala Phe Tyr Thr Ile Leu Thr Pro Met
275 280 285
Leu Asn Pro Leu Ile Tyr Ser Leu Arg Asn Lys Asp Val Ala Ala
290 295 300
Ala Leu Arg Lys Val Leu Gly Arg Cys Gly Ser Ser Gln Ser Ile
305 310 315
Arg Val Ala Thr Val Ile Arg Lys Gly
320

```

<210> 17

<211> 315

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 55036418CD1

<400> 17

```

Met Glu Thr Trp Val Asn Gln Ser Tyr Thr Asp Gly Phe Phe Leu
1 5 10 15
Leu Gly Ile Phe Ser His Ser Thr Ala Asp Leu Val Leu Phe Ser
20 25 30
Val Val Met Ala Val Phe Thr Val Ala Leu Cys Gly Asn Val Leu
35 40 45
Leu Ile Phe Leu Ile Tyr Met Asp Pro His Leu His Thr Pro Met
50 55 60
Tyr Phe Phe Leu Ser Gln Leu Ser Leu Met Asp Leu Met Leu Val
65 70 75
Cys Thr Asn Val Pro Lys Met Ala Ala Asn Phe Leu Ser Gly Arg
80 85 90
Lys Ser Ile Ser Phe Val Gly Cys Gly Ile Gln Ile Gly Leu Phe
95 100 105
Val Cys Leu Val Gly Ser Glu Gly Leu Leu Gly Leu Met Ala
110 115 120
Tyr Asp Arg Tyr Val Ala Ile Ser His Pro Leu His Tyr Pro Ile
125 130 135
Leu Met Asn Gln Arg Val Cys Leu Gln Ile Thr Gly Ser Ser Trp
140 145 150
Ala Phe Gly Ile Ile Asp Gly Leu Ile Gln Met Val Val Val Met
155 160 165
Asn Phe Pro Tyr Cys Gly Leu Arg Lys Val Asn His Phe Phe Cys
170 175 180
Glu Met Leu Ser Leu Leu Lys Leu Ala Cys Val Asp Thr Ser Leu
185 190 195
Phe Glu Lys Val Ile Phe Ala Cys Cys Val Phe Met Leu Leu Phe
200 205 210
Pro Phe Ser Ile Ile Val Ala Ser Tyr Ala His Ile Leu Gly Thr
215 220 225
Val Leu Gln Met His Ser Ala Gln Ala Trp Lys Lys Ala Leu Ala
230 235 240
Thr Cys Ser Ser His Leu Thr Ala Val Thr Leu Phe Tyr Gly Ala
245 250 255
Ala Met Phe Ile Tyr Leu Arg Pro Arg His Tyr Arg Ala Pro Ser
260 265 270
His Asp Lys Val Ala Ser Ile Phe Tyr Thr Val Leu Thr Pro Met
275 280 285
Leu Asn Pro Leu Ile Tyr Ser Leu Arg Asn Arg Glu Val Met Gly
290 295 300

```

Ala Leu Arg Lys Gly Leu Asp Arg Cys Arg Ile Gly Ser Gln His
 305 310 315

<210> 18

<211> 324

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7481701CD1

<400> 18

Met Glu Ser Pro Asn Gln Thr Thr Ile Gln Glu Phe Ile Phe Ser
 1 5 10 15
 Ala Phe Pro Tyr Ser Trp Val Lys Ser Val Val Cys Phe Val Pro
 20 25 30
 Leu Leu Phe Ile Tyr Ala Phe Ile Val Val Gly Asn Leu Val Ile
 35 40 45
 Ile Thr Val Val Gln Leu Asn Thr His Leu His Thr Pro Met Tyr
 50 55 60
 Thr Phe Ile Ser Ala Leu Ser Phe Leu Glu Ile Trp Tyr Thr Thr
 65 70 75
 Ala Thr Ile Pro Lys Met Leu Ser Ser Leu Leu Ser Glu Arg Ser
 80 85 90
 Ile Ser Phe Asn Gly Cys Leu Leu Gln Met Tyr Phe Phe His Ser
 95 100 105
 Thr Gly Ile Cys Glu Val Cys Leu Leu Thr Val Met Ala Phe Asp
 110 115 120
 His Tyr Leu Ala Ile Cys Ser Pro Leu His Tyr Pro Ser Ile Met
 125 130 135
 Thr Pro Lys Leu Cys Thr Gln Leu Thr Leu Ser Cys Cys Val Cys
 140 145 150
 Gly Phe Ile Thr Pro Val Pro Glu Ile Ala Trp Ile Ser Thr Leu
 155 160 165
 Pro Phe Cys Gly Ser Asn His Leu Glu His Ile Phe Cys Asp Phe
 170 175 180
 Leu Pro Val Leu Arg Leu Ala Cys Thr Asp Thr Arg Ala Ile Val
 185 190 195
 Met Ile Gln Val Val Asp Val Ile His Ala Val Glu Ile Ile Thr
 200 205 210
 Ala Val Met Leu Ile Phe Met Ser Tyr Asp Gly Ile Val Ala Val
 215 220 225
 Ile Leu Arg Ile His Ser Ala Gly Gly Arg Arg Thr Ala Phe Ser
 230 235 240
 Thr Cys Val Ser His Phe Ile Val Phe Ser Leu Phe Phe Gly Ser
 245 250 255
 Val Thr Leu Met Tyr Leu Arg Phe Ser Ala Thr Tyr Ser Leu Phe
 260 265 270
 Trp Asp Ile Ala Ile Ala Leu Ala Phe Ala Val Leu Ser Pro Phe
 275 280 285
 Phe Asn Pro Ile Ile Tyr Ser Leu Arg Asn Lys Glu Ile Lys Glu
 290 295 300
 Ala Ile Lys Lys His Ile Gly Gln Ala Lys Ile Phe Phe Ser Val
 305 310 315
 Arg Pro Gly Thr Ser Ser Lys Ile Phe
 320

<210> 19

<211> 312

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7481774CD1

<400> 19

```

Met Glu Pro Trp Gln His Pro Thr His Phe Ile Leu Leu Gly Phe
 1          5          10          15
Ser Asp Arg Pro His Leu Glu Arg Ile Leu Phe Val Val Ile Leu
          20          25          30
Ile Ala Tyr Leu Leu Thr Leu Val Gly Asn Thr Thr Ile Ile Leu
          35          40          45
Val Ser Arg Leu Asp Pro His Leu His Thr Pro Met Tyr Phe Phe
          50          55          60
Leu Ala His Leu Ser Phe Leu Asp Leu Ser Phe Thr Thr Ser Ser
          65          70          75
Ile Pro Gln Leu Leu Tyr Asn Leu Asn Gly Cys Asp Lys Thr Ile
          80          85          90
Ser Tyr Met Gly Cys Ala Ile Gln Leu Phe Leu Phe Leu Gly Leu
          95          100          105
Gly Gly Val Glu Cys Leu Leu Leu Ala Val Met Ala Tyr Asp Arg
          110          115          120
Cys Val Ala Ile Cys Lys Pro Leu His Tyr Met Val Ile Met Asn
          125          130          135
Pro Arg Leu Cys Arg Gly Leu Val Ser Val Thr Trp Gly Cys Gly
          140          145          150
Val Ala Asn Ser Leu Ala Met Ser Pro Val Thr Leu Arg Leu Pro
          155          160          165
Arg Cys Gly His His Glu Val Asp His Phe Leu Cys Glu Met Pro
          170          175          180
Ala Leu Ile Arg Met Ala Cys Ile Ser Thr Val Ala Ile Asp Gly
          185          190          195
Thr Val Phe Val Leu Ala Val Gly Val Val Leu Ser Pro Leu Val
          200          205          210
Phe Ile Leu Leu Ser Tyr Ser Tyr Ile Val Arg Ala Val Leu Gln
          215          220          225
Ile Arg Ser Ala Ser Gly Arg Gln Lys Ala Phe Gly Thr Cys Gly
          230          235          240
Ser His Leu Thr Val Val Ser Leu Phe Tyr Gly Asn Ile Ile Tyr
          245          250          255
Met Tyr Met Gln Pro Gly Ala Ser Ser Ser Gln Asp Gln Gly Lys
          260          265          270
Phe Leu Thr Leu Phe Tyr Asn Ile Val Thr Pro Leu Leu Asn Pro
          275          280          285
Leu Ile Tyr Thr Leu Arg Asn Arg Glu Val Lys Gly Ala Leu Gly
          290          295          300
Arg Leu Leu Leu Gly Lys Arg Glu Leu Gly Lys Glu
          305          310

```

<210> 20

<211> 1076

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7474806CB1

<400> 20

```

caaagttgaa tgccgggttg gggcagaggc tgatgccgtg ctgagggtcat gatgttatgc 60
tgtccatttt gcttccttcc aggggaagca gaagcgggag ccgtcgtgga gctctgctcc 120
tggaggggagc ctcccgggac atggagaagg tggacatgaa tacatcacag gaacaaggtc 180
tctgccagtt ctgagagaag tacaagcaag tctacctctc cctggcctac agtatcatct 240
ttatcctagg gctgccacta aatggcactg tcttgtggca ctctggggc caaaccaagc 300

```

```

gctggagctg tggcaccacc tatctggtga acctgatggt ggccgacctg ctttatgtgc 360
tattgccctt cctcatcatc acctactcac tagatgacag gtggcccttc ggggagctgc 420
tctgcaagct ggtgcacttc ctgtttcata tcaaccttta cggcagcatc ctgctgctga 480
cctgcatctc tgtgcaccag ttcttaggtg tgtggcaccg actgtgttcg ctgccctacc 540
ggacccgcag gcatgcctgg ctgggcacca gcaccacctg ggccctgggtg gtccctccagc 600
tgctgcccac actggccttc tcccacacgg actacatcaa tggccagatg atctggtatg 660
acatgaccag ccaagagaat tttgatcggc tttttgccta cggcatagtt ctgacattgt 720
ctggcctttct tccccctcc ttggtcattt tgggtgctta ttactgatg gtcaggagcc 780
tgatcaagcc agaggagAAC ctcatgagga caggcaacac agcccagacc aggtccatcc 840
ggaccatcct actggtgtgt ggcctcttca ccctctgttt tgtgcccttc catatcactc 900
gctccttcta cctcaccatc tgctttctgc tttctcagga ctgccagctc ttgatggcac 960
ccagtgtggc ctacaagata tggaggcctc tggtagtgtg gagcagctgc ctcaaccag 1020
tcctgtactt tctttcaagg ggggcaaaaa tagagtcagg ctccctccaga aactga 1076

```

<210> 21

<211> 1102

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7474840CB1

<400> 21

```

ggaggctgag gcaggagaat ggcattgaccc caggaggcag agcttgcagt gagatgagat 60
catgccactg tgctccagcc tgggcaacag agcgagactc tgtctcaaaa aaaaaaaaaa 120
acaaaaaaaa aaaccttttt tcccaagcta ccattggact tttagccaac acctttttcc 180
ttttcttcaa catcttcata ttcttcagg atcagaaatc gaagcccat gacctcatca 240
gctgtaattc ggcccttcatt catgtagtga tgttcctcac tgtggtggat gcttggcctc 300
cagatatgcc tgaatcactg cacttaggga atgagttcaa atttaagtcc ttgtcctaca 360
taaacagagt gaggatgggc ctatgtatct gtaacatctg tctcctgagt atacaccagg 420
ccaacaccat cagccccaac aacttctgtt tggcaaggct taaacagaaa ttcacaaata 480
acattatcat gtcattcttt ttttctttt ttttttggtc catcaatttg tctttcagtt 540
ataacatagt attctttact gtggcttctt ctaatgtgac ccagaacagt ctacctaaag 600
gcagcaatac tgttcacttt ctcccatga agtccttcat gagaaaagta ttttttactc 660
tgacattatc cagggatgtc ttcatatag gaattacact gcattcaatt gcacacatgg 720
tgatccttgt gtccaggcat gagacgcaat ctacagacct tcacagcatc agcatctctc 780
cacaagcctt cccagagaaa agggctgtc agaccatccc gctgttagtg agctactgtc 840
tggctcatgt ctgggtggac ctcatcatct catctcttc aacctgtctg tggacgtgta 900
accagctctt cctgagtatg cagaaccttg tggcgatgt ctatgccact gttgttctac 960
tggaaacaaat cagctctgat aaaaatatag ttgacattct ccaaaatatg caaagtgtca 1020
taaagcttta acaagttggc gatggaaaac atttctaaaa aatagtcttc tcctatagtt 1080
caattgttca agtagccctg ga 1102

```

<210> 22

<211> 2529

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7475092CB1

<400> 22

```

ggctccgggtg tttcccgccg ttcatgcagc gaaaagagaa agcaaaatgc cctcaggagg 60
ctccagcccg cgcgagccc tccacgcccg gcgggggcag cggaggcgga ggcgccgtcg 120
ctgcagcctc aggcgcgcg gtgcgggct cgtgcagtt ggcgtgagc gtccctgcagc 180
ccctgctcta cgcgcgctg ttgcctttg cctacctgca gctgtggcgg ctgctcctgt 240
accgcgagcg gcggctgagt taccagagcc tctgcctctt cctctgtctc ctgtgggcag 300
cgctcaggac caccctcttc tccgcgcct tctcgtcag cggctccctg cccttgctcc 360
ggcgcgccgc tcacctgcac ttcttcccc actggctgt ctactgtctc ccctcctgtc 420
tccagttctc cagctctgt ctctcaacc tctacctggc ggaggttata tgtaaagtca 480
gatgtgccac tgaacttgac agacacaaaa ttctactgca tttgggcttt ataattggca 540

```

```

gcctgctctt ttttagtggtg aacttgactt gcgcaatgct agttcatgga gatgtcccag 600
aaaatcagtt gaagtggact gtgtttgttc gagcattaat taatgatagc ctgtttattc 660
tttgtgccat ctcttttagtg ttttacatat gcaaaattac aaaaatgtca tcagctaattg 720
tctacctga atcaaagggg atgtctctgt gccagactgt cgctcgtgggc tctgtagtca 780
ttcttctgta ctctccaga gcttgttata atttgggtgg ggccaccata tctcaggata 840
cattagaag tccatttaat tatggctggg ataactcttc agataaggct catgtagaag 900
acataagtgg agaagagtat atagtatttg gaatggctct ctttctgtgg gaacatgtgc 960
cagcatgggc ggtgggtactg ttttccggg cacagagatt aaaccagaat ttggcacctg 1020
ctggcatgat aaatagtcac agttatagtt ccagagctta ctttttcgac aatccaagac 1080
gatatgatag tgatgatgac ctgccaagac tgggaagtgc aagagaagga agtttaccac 1140
attcgcaaag tttgggctgg tatggcacca tgactgggtg tggcagcagc agttacacag 1200
tcactcccca cctgaatgga cctatgacag tactgctcc tttgctcttt actttagtga 1260
atttagattt gaacaatcat catagcttat atgtgacacc acaaaactga cagcatcacc 1320
aagtcagatg tcttgagttg ttttccataa atgtgtatat tcaatgtgtt taaattccat 1380
ctacataaac attccattat ctgttgcaac tgaaaacaaa atctggaagt gtggctgtgt 1440
ttggtaaata acacagctat tatttttgac ctcttcatag taaaatgaag taaaatggaa 1500
agtttggagt aggagaaaag agagattaga tcttaaggca cttgatggcc tccaaaaatc 1560
ctgactttgg aacatcaaat gcatatgtgc acttttatct ttgttctgag tccactgcag 1620
cccaaagtc atatgccaat gttcacactg aaatactgta ttgtacacca aactggaagg 1680
caattttcct atgaaaatca aagccggtat attcattggg atgctctata cagatatctt 1740
aataaaaatt ttatagtgtg aacagtgcac agagttaagg cataaaaatg tatcattctt 1800
tataaaaatc tactgaaaat gtgtaatcat tgaagacagt tcttttaagc atgattttaa 1860
aatagcaact gaaattcaat catttttaac aaatgatggg agtaatccat tagttatggc 1920
cagcagtgtt ctttgagag ccacaataat ttcaagagga aaatatacca gtgaaaattg 1980
tgtggctatt ttgagtagaa ttggtcagtt gattattttg tgtaattgag atatatgtag 2040
tagtttaagc atgattcttg aagaaagcaa tagtgacttt tgcataggga gattttggta 2100
gaaacttctt gggactaaac aagtttagag atgcatttaa gaattattca caaaatgtgt 2160
aattctaaat taaaacataa atatattttc aaaagcattt gatttctctg aagcatgata 2220
tagctggtct tacctagtga atcaggattg tcctcaggta aatgaaatca tgatacatta 2280
ttgcagtga ctcaagtga atactttgta agacatataa ttccatgatg tttcacattt 2340
ttatatctta tatatgggaa aagccaaatt aaattgaatt cagattaatt ccagcattag 2400
actaaatgag caaacttaag taaatgtaca aactaggtaa gtataaaacc acaggttaac 2460
aatattggag tactttttaga attacattaa aactgtctta aatgtcctat cccaaatcta 2520
aaaaaaaaa 2529

```

<210> 23

<211> 1847

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7341260CB1

<400> 23

```

gggggaggca ggcctcccag ttgctgcagt ttggaatatg tcagggtccca cctcccaga 60
ggcggggcca gggctgagtc ctgccagcct catttctcta tccctctgag aaccagacg 120
ggcagagcct gggtaggaga gcctggcccc gctgtcccca ctgggtggag acaccatgca 180
cttgggtccac ttgtgctctt cagccaggac accagacatg gtccaaaccg ctgcagggtc 240
ggctgcagca actcctgac actcaggaag gcccaggctg ggcaggcaat acctgctccc 300
aacagccatg catgccggct gccgtccag gaactccctg tccccaggac caagatgacg 360
cccaacagca ctggcgaggt gccagcccc attcccaagg gggctttggg gctctccctg 420
ggcctggcaa gcctcatcat caccgcgaac ctgctcctag ccttgggcat cgctgggac 480
cgccgctgc gcagccacc tgctggctgc ttcttctga gcctactgct ggctgggctg 540
ctcacgggtc tggcattgcc cacattgccca gggctgtgga accagagtcg ccgggggttac 600
tggctcctgcc tectctgcta cttggctccc aacttctcct tctcttccct gcttgccaac 660
ctcttgctgg tgcacgggga gcgctacatg gcagtcctga ggccactcca gcccctggg 720
agcattcggc ttggcctgct cctcacctgg gctgttcccc tgctctttgc cagtctgccc 780
gctctggggg ggaaccactg gaccctgggt gccaaactgca gctcccaggc tatcttccca 840
gccccctacc tgtacctga agtctatggg ctctctgctg ccgccgtggg tgctgctgcc 900
ttctctctg tccgctgct ggccactgcc caccgccagc tgcaggacat ctgccggctg 960
gagcgggcag tgtgcgcga tgagccctcc gccctggccc gggcccttac ctggaggcag 1020
gcaagggcac aggcctggagc catgctgctc ttccgggctgt gctggggggc ctacgtggcc 1080

```

```

acactgctcc tctcagtcct ggccatagag cagcgcccg cactggggcc tgggacactg 1140
ttgtccctcc tctccctagg aagtgccagt gcagcggcag tgcccgtagc catggggctg 1200
ggcgatcagc gctacacagc cccctggagg gcagccgccc aaaggtgcct gcaggggctg 1260
tggggaagag cctcccggga cagtcccggc cccagcattg cctaccaccc aagcagccaa 1320
agcagtgtcg acctggactt gaactaaagg aagggcctct gctgactcct accagagcat 1380
ccgtccagct cagccatcca gcctgtctct actgggcccc acttctcttg atcagagacc 1440
ctgcccctgt ttgaccccg actgactgaa taaagctcct ctggccgtta aaaaaaaaaa 1500
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaacaaacag 1560
aaaaaaaaaa aaaagaacac acaaagaaca cagaacaaac caagcagcac accacacaca 1620
aaaacaatgc acacacagaa caagacacaa tcagagacag agagcacaca gcacggaccc 1680
cagccacgcc cccagcactg accaccacga cccgacacag aaacgaacac tgaagactca 1740
acgcacaaaa cgacaaccag accacaagcc aaccgcctca cgccccagca acgaacacac 1800
atacaaaacc aaaccgagac aaccacata cagccaaaca aaccaca 1847

```

<210> 24

<211> 2031

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7473911CB1

<400> 24

```

atgaataaaa acaacaaacc ttccagtttc atagccataa gaaatgctgc tttctctgaa 60
gtcggcattg ggatctctgc caatgccatg ctccctctct tccacatcct cactgtgcct 120
ctcaagcaca ggaccaagcc cgctgacctg atcgtttctc atgtggctct aatccatctc 180
atattgctgc taccacaga gttcatagct acagatatct ttgggtctca ggattcagag 240
gatgacatca aacataagtc agttatctac aggcgtaaca ggcagtccca gcattttcac 300
agcaccaacc tttctccaaa agcaccccca gaaaaaatgg ccacgcagac cattcttctg 360
ctcgtgagtt gctttgtgat tgtgtatggt ttggactgtg ttgtcgctc ctgctcagga 420
ctgggtgtgga acagtgatcc agtccgtcat cgagtccaga tgctgggtgga caatggctat 480
gccaccatca gtccttcagt gctaccagg ctgactgccc caaacgagtg gagagccagt 540
gtgtacctga atgacagctt gaacaaatgc agcaacggac ggctgctctg tgtagacagg 600
gggcttgatg aggggccccg gtcctgcccc aagtgtctct agtcagagac cgacgaggat 660
tacatcgctc tcagggtctc gctgaggagg gacgaaccca aggacggggg cagtgtgggg 720
aatgcagccc tgggtgtctc cgaggcctct gcagaagagg aagaggagcg tgaggaggga 780
ggcgaggcat gtgacctgga gaggacagga gctgggtggg agcagggtga ccttggtgaa 840
ctacctgacc atgaggagaa aagcaaccag aaagtggcag ctgccaccct ggaggaccgc 900
acacaggatg agcctgtgga ggagagctgc cagatcgctc ttttccagaa caactgcatg 960
gacaactttg tgacttccct cacaggaagc ccctacgagt tcttcccaac caagagcacc 1020
tctttttgca gggagagctg ttctcctttt tctgagtcag tgaaaagctt agaatcagag 1080
caggcaccaa agttggggct gtgtgcggag gaggacccc tggttggggc tttgtgtggc 1140
cagcatggac ccttgcaaga tggagtggcg gagggtccca cagcccctga tgtggtggtc 1200
ctgccgaagg agggaggaga ggaggaggtc attgtggatg acatgctggc caaccctat 1260
gtgatggggg atgaggggga ggaggaggag gaggagtctg tggatgacac actggccaac 1320
ccctatgtga tgggagtggg cctgccagga agaggagggg agggaggaga ggaggaggag 1380
gtcgtggatg acacgctggc cagcctctat aagatgggag aagaacatcg acacaagggc 1440
ctggccccac tctgggaagg tggccagaaa ccgtcccaga aactgcccc aaagaaacca 1500
gatctgaggg aggttctctc gcccttgcca tcggagggtc cgcagaggag gcaggaaaga 1560
gctgttgtca ctgaaggagg gcccttgga gccagcagg ccttgccagc aaagcccagg 1620
gccttcactt tataccctcg gtcgttctcc gtggaaggcc aagagattcc tgtttccatc 1680
tctgtgtact gggagccaga agggtcgggg ttagatgacc acaggataaa gaggaagag 1740
gaacatctct ctgtcgtgtc tgggagtctc tcccagagaa accaccttcc atccagcggc 1800
acctccacgc cttcttccat ggtcgacatc ccacctcct tcgacctggc ctgcatcacc 1860
aagaagccca tcacaaagag ctctccctct ctctctgatg acagcgactc cccggacaag 1920
tacaagaaga agaagtcata ctttaagcgg ttcttgccgc tgatgtttaa caagatggag 1980
aggccaggca cgatggctca tgcctgtcat ccagcactt tgggaagctg a 2031

```

<210> 25

<211> 1130

<212> DNA

<213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7474767CB1

<400> 25
 ggggcgcgct catggagcac acgcacgccc acctcgcagc caacagctcg ctgtcttggt 60
 ggtcccccg ctcggcctgc ggcttgggtt tegtgcctgt ggtctactac agcctcttgc 120
 tgtgcctcgg tttaccagca aatatcttga cagtgatcat cctctcccag ctggtggcaa 180
 gaagacagaa gtccctcctac aactatctct tggcactcgc tegtgcgcag atcttgggtcc 240
 tctttttcat agtgtttgtg gacttctctgt tgggaagatt catcttgaac atgcagatgc 300
 ctcagggtccc cgacaagatc atagaagtgc tgggaattct atccatccac acctccatat 360
 ggattactgt accgttaacc attgacaggt atatcgctgt ctgccaccgc ctcaagtacc 420
 acacggtctc ataccagcc cgcacccgga aagtcattgt aagtgtttac atcacctgct 480
 tcctgaccag catcccctat tactggtggc ccaacatctg gactgaagac tacatcagca 540
 cctctgtgca tcacgtcctc atctggatcc actgcttcac cgtctacctg gtgccctgct 600
 ccattcttct catcttgaac tcaatcattg tgtacaagct caggaggaag agcaattttc 660
 gtctccgtgg ctactccacg gggaagacca ccgccatctt gttcaccatt acctccatct 720
 ttgccacact ttgggcccc cgcacatca tgattcttta ccacctctat ggggcgccca 780
 tccagaaccg ctggctggta cacatcatgt ccgacattgc caacatgcta gcccttctga 840
 acacagccat caacttcttc ctctactgct tcacagcaa gcggttccgc accatggcag 900
 ccgcccagct caaggcttct tccaagtgc agaagcaacc tgtacagttc tacaccaatc 960
 ataacttttc cataacaagt agcccctgga tctcgccggc aaactcacac tgcatcaaga 1020
 tegtgtgtga ccagtatgac aaaaatggaa aacctataaa aagtcgtaat gacagcaaaa 1080
 gctcctacca gtttgaagat gccattggag cttgtgtcat catcctgtga 1130

<210> 26
 <211> 1202
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7475815CB1

<400> 26
 caggttctgc aatacaattg gaaacaactc attgtctctg cctatgaaga agtagaacta 60
 tggtaaaaaat aagaaaatgc cttcccaata ataggaggtt gtactatgga agtaatatgg 120
 agggccagca atgggaatca tagtccagcc atgatggcat agtcatggga gaagagagag 180
 ggtggggatg gcttccagga ggggtgtgaag aggggggtcca gtactgaagg gggaaaagat 240
 gcatcagaat taattaatgt attttgatga tggcaatagt gttgggttgag attggtgaag 300
 gtagtaatat ttgtgatatt tttgttgctt ttctccctag acatthaacta tgtgcttatt 360
 ttccccataa gatgaataaa aacaacaac cttccagttt catagccata agaaatgctg 420
 ctttctctga agtcggcatt gggatctctg ccaatgccat gctccttctc ttccacatcc 480
 tcacgtgcct tctcaagcac aggaccaagc ccgctgacct gatcgtttgt catgtggctc 540
 taatccatat catattgctg ctaccacacag agttcatagc tacagatatt tttgggtctc 600
 aggattcaga ggatgacatc aaacataagt cagttatcta caggtagagg ttgatgagag 660
 gcctctccat ttccaccacc tgcctgctga gtatcctccc ggccatcacc tgcagcccca 720
 gaagctcctg tttggcagtg ttcaaagatt ctcacatcac caaccaggtt gctttctctt 780
 ccgtcttcca catatccatt agtgacagct tcttagtctc cactcttccc atcaaaaatc 840
 tggcctcaaa tagccttaca tttgtcactc aatcctgctc tgcctgggatc ggctcacggc 900
 cccctccag tggatacatg gtgattctct tgtccaggcg taacaggcag tcccagcatt 960
 ttcacagcac caacctttct ccaaaagcac cccagaaaaa aatggccacg cagaccattc 1020
 ttctgctcgt gagttgcttt gtgattgtgt atgttttggg ctgtgttgct gcctcctgct 1080
 caggactggg gtggaacagt gatccagtc gtcacagagt ccagatgctg gtggacaatg 1140
 gctatgccac catcagtcct tcagtgttag tcagtactga aaaatgaatg atcaaaagtct 1200
 ga 1202

<210> 27
 <211> 2079
 <212> DNA
 <213> Homo sapiens

<220>

<221> misc_feature
 <223> Incyte ID No: 60263275CB1

<400> 27

```

tacggcacag tagagagctt ccagggctgg ctggcggtgg ataccctgtac cacagaaatg 60
cagggaccat tgcttcttcc aggcctctgc tttctgctga gcctcttttg agctgtgact 120
cagaaaacca aaaacattaa tgaatgtaca ccaccctata gtgtatatatg tggatttaac 180
gctgtgtgtt acaatgtcga aggaagtttc tactgtcaat gtgtcccagg atatagactg 240
cattctggga atgaacaatt cagtaattcc aatgagaaca cctgtcagga caccacctcc 300
tcaaagacaa cccagggcag gaaagagctg caaaagattg tggacaaatt tgagtcactt 360
ctcaccaatc agactttatg gagaacagaa gggagacaag aaatctcatc cacagctacc 420
actattctcc gggatgtgga atcgaaagt ctgaaaactg ccttgaaaga tccagaacaa 480
aaagtcctga aaatccaaaa cgatagtgtg gctattgaaa ctcaagcgat tacagacaat 540
tgctctgaag aaagaaaagac attcaacttg aacgtccaaa tgaactcaat ggacatccgt 600
tgcaagtaca tcatccaggg agacacacaa ggtcccagtg ccattgcctt tatctcatat 660
tcttctcttg gaaacatcat aaatgcaact ttttttgaag agatggataa gaaagatcaa 720
gtgtatctga actctcaggt tgtgagtgt gctattggac caaaaaggaa cgtgtctctc 780
tccaagtctg tgacgctgac tttccagcac gtgaagatga ccccagtag caaaaaggtc 840
ttctgtgtct actggaagag cacagggcag ggcagccagt ggtccaggga tggctgcttc 900
ctgatacacg tgaacaagag tcacaccatg tgaattgca gtcacctgtc cagcttcgct 960
gtcctgatgg ccctgaccag ccaggaggag gatcccgtag tgactgtcat cacctacgtg 1020
gggctgagcg tctctctgct gtgcctcttc ctggcgccgc tcaactttct cctgtgtaaa 1080
gccatccaga acaccagcac ctcaactgcat ctgcagctct cgctctgcct ctccctggcc 1140
cacctctctt tctctgtggg gattgatcga actgaaccca aggtgctgtg ctccatcatc 1200
gcgggtgctt tgcactatct ctacctggcc gccttcacct ggatgctgct ggaggggtgtg 1260
cacctcttcc tcactgcacg gaacctgaca gtggtcaact actcaagcat caatagactc 1320
atgaagtgga tcatgttccc agtcggctat ggcgttcccg ctgtgactgt ggccatttct 1380
gcagcctcct ggcctcacct ttatggaact gctgatcgat gctggctcca cctggaccag 1440
ggattcatgt ggagtttctt tggcccagtc tgtgccattt tctctgcaa tttagtattg 1500
tttatcttgg tcttttggat tttgaaaaga aaactttcct ccctcaatag tgaagtgtca 1560
accatccaga acacaaggat gctggctttc aaagcaacag ctcaactctt catcctgggc 1620
tgcacatggg gtctgggctt gctacagggt ggtccagctg cccaggctcat ggcctacctc 1680
ttcaccatca tcaacagcct ccaaggcttc ttcattctct tgggtctact cctcctcagc 1740
cagcaggtcc agaaaacaata tcaaaagtgg tttagagaga tcgtaaaatc aaaatctgag 1800
tctgagacat acacactttc cagcaagatg ggtcctgact caaaacccag tgagggggat 1860
gtttttccag gacaagtga gaaaaatat taaaactaga atattcaact ccatatggaa 1920
aatcatatcc atggatctct ttggcattat gaagaatgaa gctaaggaaa aggggaattca 1980
ttaaacatat catccttggg gaggaagtaa tcaaccttta cttcccaaac tgtttgttct 2040
ccacaatagg tctcaacaaa tgtgtggtaa attgcatta 2079

```

<210> 28
 <211> 5324
 <212> DNA
 <213> Homo sapiens

<220>

<221> misc_feature
 <223> Incyte ID No: 60203310CB1

<400> 28

```

ggcggcagca gcagcagcaa gaggaggaga agcagcccca gctatgactg ccgcatgtta 60
atagctgccg ctgggtccct cggctgctgc tggagacaga gcctgactcc gaagttgtgc 120
aactgtggac tgggagagac atttgaacct tcttttcttt tcgctccctt tttgcccct 180
tgggggtgtg gaagcgagga acgtaaagga aggcgaacat ttggctctct ttttccctcc 240
cctttctccg tggctgtgta gcggaagaaa gggagagag actttttgtt gttgtttcct 300
tgactggggt ctccaccctc ctgctgcttt cctcgcgctt cgattctcgt tatttgccg 360
gtgtgggttg ggggtgtctg acaggggccc gccggtcttt tgccccgggc tcaatggctg 420
gattgtggaa actgcacccg ccttcagggt gttgagcaac tgatgggacg atctcagga 480
ccggcgttta cgaaagggtt cagatttggg atattgggtt ttctgttttg gagaaattat 540
tcttttcttt ttttaattga agaaaaatca tcagtcttgg aatacagaag agaaactaga 600
aatatacgta ttttgtttca catttgaaca gtcattcttg aggaatactc catacctgag 660
tagacagcca tgtggccatc gcagctacta atttctatga tgctcttagc tccaataatt 720
catgctttca gccgtgcccc aattccaatg gctgtggtcc gcagagagct atcctgtgag 780

```

agctatccta	tagagcttcg	ctgtccagga	acagacgtca	tcatgataga	aagtgccaac	840
tatggcagga	ctgatgacaa	aatttgtgac	tctgaccctg	ctcagatgga	gaatatccga	900
tgttatctgc	cagatgccta	taagattatg	tctcaaagat	gcaataacag	aaccagtggt	960
gcagtgggtg	caggtcctga	tgtttttcca	gacccgtgtc	caggaacctta	taaatacctt	1020
gaagtgcagt	atgaatgtgt	cccttacaaa	gtggaacaaa	aagtttttct	ttgtcctgga	1080
ctactaaaag	gagtatacca	gagtgaacat	ttgtttgagt	ccgaccacca	atctggggcg	1140
tgggtgcaaag	accctctgca	ggcatctgac	aagatttatt	atatgccctg	gactccctac	1200
agaactgata	ccctgactga	gtattcatcc	aaggatgact	tcattgctgg	aagaccaact	1260
acaacctaca	agctccctca	taggggtggat	ggcacaggat	ttgtagtgtta	tgatggagct	1320
ttgttcttca	acaaagagcg	caccaggaac	atagtaaagt	ttgatttgcg	gactaggata	1380
aagagtggag	aggctatcat	agcaaagtcc	aattaccatg	atacctcccc	ttaccgatgg	1440
ggaggcaaat	ctgacataga	cctggcagta	gatgagaatg	ggctatgggt	aatctatgca	1500
acagaacaaa	acaatggtaa	aattgtcatt	agtcaattga	acccttacac	cctacggatc	1560
gaagggaacat	gggatactgc	atatgataaa	aggtcagctt	ccaatgcctt	tatgatttgt	1620
ggaattctgt	atgtggtcaa	atctgtatat	gaggatgatg	acaatgaggc	tactggaaat	1680
aagattgact	acattttacaa	cactgaccac	agcaaggata	gtttggtgga	tgtacccttt	1740
cctaattcat	accagtacat	tgcagctgtg	gattacaacc	ccagggacaa	cctactttat	1800
gtatggaata	actatcacgt	cgtgaaatat	tctttggatt	ttggacctct	ggatagtaga	1860
tcaggggcagg	acatcatggt	acaagtttca	tacatttctc	cgccaattca	ccttgactct	1920
gagctagaaa	gacctctgtg	taaagatatc	tctaccacag	gacctcttgg	catgggaagc	1980
actaccacca	gtaccaccct	tcggaccaca	actttgagcc	cagggaaggag	taccaccccc	2040
tcagtgtcag	gaagaagaaa	ccggagtact	agtaccccat	ctccagctgt	cgagggtact	2100
gatgacatga	ccacacacct	tccatcagca	tcgtcccaaa	tccagctctc	cgaagagagc	2160
tgtgaggctg	tgggaagccc	agaaatcatg	tggtttaaga	ctcgtcaagg	acagatagca	2220
aagcagccat	gccctgcagg	aactataggt	gtatcaactt	atctatgcct	tgctcctgat	2280
ggaatttggg	atccccaagg	tccagatctc	agcaactggt	cttctccttg	ggtcaatcat	2340
ataacacaga	agttgaaatc	tggtgaaaca	gctgccaaac	ttgctagaga	gctggctgaa	2400
cagacaagaa	atcacttgaa	tgctggggac	atcacctact	ctgtccgggc	catggaccag	2460
ctggtaggcc	tcttagatgt	acagcttcgg	aacttgaccc	cagggtggaa	agatagtgct	2520
gcccggagtt	tgaacaagct	tcagaaaaga	gagcgtctct	gcagagccta	tgtccaggca	2580
atggtcgaga	cagttaacaa	cctccttcag	ccacaagctt	tgaatgcatg	gagagacctg	2640
actacgagtg	atcagctgcg	tgccggccacc	atgttgcttc	atactgtgga	ggaaagtgtc	2700
tttgtgctgg	ctgataacct	tttgaagact	gacattgtca	gggagaacac	agacaatatt	2760
aaattgggaag	ttgcaagact	gagcacagaa	ggaaacttag	aagacctaaa	atttccagaa	2820
aacatggggc	atggaagcac	tatccagctg	tctgcaaata	ccttaaagca	aaatggccga	2880
aatggagaga	tcagagtggc	ctttgtcctg	tataacaact	tgggtcctta	tttatccacg	2940
gagaatgccca	gtatgaagtt	gggaacggaa	gctttgtcca	caaatcatte	tgttattgtc	3000
aattcccctg	ttattacggc	agcaataaac	aaagagtcca	gtaacaaggt	ttatttggct	3060
gatctgtggt	tattttactgt	taaacatatc	aagcagtcag	aggaaaattt	caaccttaac	3120
tgttcatttt	ggagctactc	caagcgtaca	atgacaggtt	attggtcaac	acaaggtgt	3180
cggtccttga	caacaaataa	gacacatact	acatgctctt	gtaaccacct	aacaaatttt	3240
gcagtactga	tggcacatgt	ggaagttaag	cacagtgatg	cggtccatga	cctccttctg	3300
gatgtgatca	cgtgggttgg	aattttgtct	tcccttggtt	gtctcctgat	ttgcatcttc	3360
acatttttct	ttttccggcg	gctccagagt	gcacgttaaca	ccatccacaa	gaacctctgc	3420
atcagctctc	ttgtagcaga	gctgctcttc	ctgattggga	tcaaccgaac	tgaccaaccg	3480
attgcctgtg	ctgttttctc	tgccctgtta	catttcttct	tcttggtgct	cttcacctgg	3540
atgttccttg	aggggggtgca	gcttttatatc	atgctggtgg	aggtttttga	gagtgaacat	3600
tcacgtagga	aatactttta	tctgggtcggc	tatgggatgc	ctgcactcat	tgtggctgtg	3660
tcagctgcag	tagactacag	gagttatgga	acagataaag	tatgttggct	ccgacttgac	3720
acctacttca	tttggaagttt	tataggacca	gcaactttga	taattatgct	taatgtaatc	3780
ttccttggga	ttgctttata	taaaatggtt	catcatactg	ctatactgaa	acctgaatca	3840
ggctgtcttg	ataacatcaa	ctatgaggat	aacagaccct	tcatacaagt	atgggttata	3900
ggtgcaatag	ctcttctctg	cctattagga	ttgacctggg	cctttggact	catgtatatt	3960
aatgaaagca	cagtcacatc	ggcctatctc	ttcaccattt	tcaattctct	acagggaatg	4020
tttatattta	ttttccattg	tgctctacag	aagaaggtag	gaaaagagta	tgggaaatgc	4080
ctgccaacac	attgctgtag	tggcaaaaagt	acagagagtt	ccattgggtc	agggaaaaca	4140
tctggtttct	gaactcctgg	acgctactcc	acaggctcac	agagccgaat	ccgtagaatg	4200
tggaatgaca	cggttcgaaa	gcagtcagag	tcttccctta	ttactggaga	cataaacagt	4260
tcagcgtcac	tcaacagaga	ggggcttctg	aacaatgcca	gggatacaag	tgtcatggat	4320
actctaccac	tgaatggtaa	ccatggcaat	agttacagca	ttgccagcgg	cgaataacctg	4380
agcaactgtg	tgcaaatcat	agaccgtggc	tataaccata	acgagaccgc	cctagagaaa	4440
aagattctga	aggaaactcac	ttccaactat	atcccttctt	acctgaacaa	ccatgagcgc	4500
tccagtgaac	agaacaggaa	tctgatgaac	aagctggtga	ataaccttgg	cagtgggaag	4560

```

gaagatgatg ccattgtcct ggatgatgcc acctcgttta accacgagga gagtttgggc 4620
ctggaactca ttcatgagga atctgatgct cctttgtctgc cccaagagt atactccacc 4680
gagaaccacc agccacacca ttataccaga aggcggatcc cccaagacca cagtggagagc 4740
tttttccctt tgctaaccac cgagcacaca gaagatctcc agtcacccca tagagactct 4800
ctctatacca gcatgccgac actggctggt gtggccgcca cagagagtgt taccaccagc 4860
acccagaccg aacccccacc ggccaaatgt ggtgatgccg aagatgttta ctacaaaagc 4920
atgccaaaac taggctccag aaaccacgtc catcagctgc atacttacta ccagctaggt 4980
cgcggcagca gtgatggatt tatagttcct ccaaaacaag atgggacccc tcccagggga 5040
agttcaaaaag gaccggctca tttggtcact agtctataga agatgacaca gaaattggaa 5100
ccaacaaaac tgctaaccac ttgttgactg ttctgagttg atataagcag tggtaataat 5160
gtgtgtactc ctaaattcttt atgctgtcct ctaaaagaca acacaaactc tcagactttt 5220
ttttttcaac tgggatttaa ggtcagccca ggggagaaa ataatgcta aaattccctt 5280
gtaccccatc ctttcttgtc ctttccccct tcagatggag actt 5324

```

<210> 29

<211> 1962

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7477349CB1

<400> 29

```

atggatccca gcgttgttag caatgagtat tatgatgttg cccatggagc aaaagatcca 60
gtgggtccca cttccctgca ggacatcact gctgtcctgg gtacagaagc atatactgag 120
gaagacaaat caatgggtgc ccatgcacag aaaagccagc attcttgtct cagccattcc 180
aggtggctga ggtctccaca ggtcacaggg ggaagctggg acctccgaat aaggccatcc 240
aaggactcca gcagtttccg ccaggctcag tgtctgcgta aggatcctgg ggcaaacaac 300
cacttgagga gccaaaggggt gagaggtaca gctggcgatg ctgacaggga gctgcgggga 360
ccctcagaaa aagccacagc tggccagcca cgagtgacct tgctgcccac gcccacgtc 420
agcgggctga gccaggagt ttgaaagccac tggccagaga tcgcagagag gtccccgtgt 480
gtggctggcg tcatccctgt catctactac agtgtcctgc tgggcttggg gctgcctgtc 540
agcctcctga cgcagtggt cctggcgcg cttgccacca ggaccaggag gccctcctac 600
tactaccttc tggcgctcac agcctcggat atcatcatcc aggtggctcat cgtgttcgcg 660
ggcttccctc tgcaggagc agtgtggcc cgccagggtg cccaggctgt ggtgcgcagc 720
gccaacatcc tggagtttgc tgccaaccac gcctcagtc ggatcgccat cctgctcacg 780
gttgaccgct acaactgcct gtgccacccc ctgcaccatc gggccgcctc gtccccaggc 840
cggacccgcc gggccattgc tgctgtcctg agtgtgtccc tgttgaccgg catccccttc 900
tactggtggc tggacatgtg gagagacacc gactcaccga gaacactgga cgaggtcctc 960
aagtgggctc actgtctcac tgtctatttc atcccttgtg gcgtgttccct ggtcaccaac 1020
tcggccatca tcaccgggt acggaggagg ggccggagtg ggctgcagcc ccgggtgggc 1080
aagagcacag ccactcctct gggcatcacc acaactgttc cctcctgtg ggcgccccgg 1140
gtcttcgtca tgctctacca catgtacgtg gcccctgtcc accgggactg gagggtccac 1200
ctggccttgg atgtggccaa catggtggcc atgtccaca cggcagccaa cttcggcctc 1260
tactgctttg tcagcaagac tttccgggcc actgtccgac aggtcatcca cgatgcctac 1320
ctgccctgca ctttggcatc acagccagag ggcattggcg cgaagcctgt gatggagcct 1380
ccgggactcc ccacaggggc agaagtgtag aggagggggc ccagctaggg agctcaggg 1440
ggctcatggc cacatgtact ggggcctttg aggttgtacc caaaacacgt ttatcaacag 1500
cttgctttcc ttgggtgggg gtggaggctc ctcccttggg tgtggctccc aggtagagag 1560
gaggacaact tagccagctc ttatgtttgc ttcaccagca atccctattt cctgggaaga 1620
tgaaagggca ctgccaggca caggctaata gcatcagtgc tgtgggcatt cctttgcggg 1680
gggcattttg cctggctcat cgtgaatgcc agattaatgt tggttgaatg gatagaaaaa 1740
cggcctctca ttttcgtaac tgaggcagga gaatcgcttg aaccaggag acggaggttg 1800
cagcgagctg agatcgcgcc atagaaacac catggaactc caacctgggc aacaagagt 1860
aaacttcgac tcaaaaaaaaa aagagaaaaa acacattagg taacagtttc tttttagcat 1920
ttgtgtaacc ttaataaaaa taaagtgata atcaaaaaaa aa 1962

```

<210> 30

<211> 1558

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 55002225CB1

<400> 30

```

atttttattcg cgaaggcacc ccacgctcct agaaaagagc acgacgcacc cgatgctcgg 60
attggatgaa gtggcacaagc tttaatccct ggaaagtcca cgaacaatga atccatttca 120
tgcattcttg tggaaacacct ctgccgaact tttaaacaaa tcctggaata aagagtttgc 180
ttatcaaaact gccagtgtgg tagatacagt catcctccct tccatgattg ggattatctg 240
ttcaacaggg ctggttggca acatcctcat tgtattcact ataataagat ccaggaaaaa 300
aacagtcctt gacatctata tctgcaacct ggctgtggct gatttgggtc acatagttgg 360
aatgcctttt cttattcacc aatgggcccg agggggagag tgggtgtttg gggggcctct 420
ctgcaccatc atcacatccc tggatacttg taaccaattt gcctgtagtg ccatcatgac 480
tgtaatgagt gtggacaggt actttgccct cgtccaacca tttcgactga cacgttggag 540
aacaaggtac aagaccatcc ggatcaattt gggcctttgg gcagcttcct ttatcctggc 600
attgcctgtc tgggtctact cgaaggtcat caaatttaaa gacggtgttg agagttgtgc 660
ttttgatttg acatcccttg acgatgtact ctggtataca ctttatttga cgataacaac 720
tttttttttc cctctaccct tgattttggt gtgctatatt ttaattttat gctatacttg 780
ggagatgtat caacagaata aggatgccag atgctgcaat cccagtgtac caaaacagag 840
agtgtatgaag ttgacaaaaga tgggtctggt gctggtggta gtctttatcc tgagtgtgc 900
cccttatcat gtgatacaac tgggtgaactt acagatggaa cagccacac tggccttcta 960
tgtgggttat tacctctcca tctgtctcag ctatgccagc agcagcatta acccttttct 1020
ctacatcctg ctgagtggaa gcctcaaat ccaaagaaga gcgactgaga aggaaatcaa 1080
caatatggga aacactctga aatcacactt ttaggaaagt acatggatca ccatgagtct 1140
agacatgatt gtctatctta ctggtattat tagaaagggc aggtgtaccg atatgtttat 1200
gccattctt cttgtgtact tgtgactctt agcagcatgg aagagaagtg taaccatgca 1260
aatacaatga gcttaatatg ctaactttag caagatgtaa aatgttgatc tatattgttg 1320
gtagggaatg ggatagtctg agataccagc gcttcagat ggtgtatatt atttcagcat 1380
attataaact agtcactaat gaaaatggcc atccatgacc attgactcaa aactcaccaa 1440
ggaacctgac cttgccctcc aactgcggc ctactgtaa cagtttctc aaggttctta 1500
ggagggtatc accttagagt gaagtctaaa atttggctat tttttatcta ttaaaaaa 1558

```

<210> 31

<211> 2304

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7475686CB1

<400> 31

```

atgctggctgg gacctgtccc agcccgggcg cgcgccctct tgtcttgggt tagggggctg 60
gaaagccgag gaggggagtg gaccaaagtc attgttcagc tgggtcatct ccttgctacc 120
cagcatcccg cggcgccccc atgtggagtc gtttccagcg ccctgggtcat gactcaaca 180
gatgtctgtc tagcccccac tatgcaacag gcactggact gggcagcagg aatttggttt 240
acaggaagat taggactcag agagcataaa tcaactggccc aggttgactc agtctgtcca 300
tgtgaaagtg aacttgggtg tttccaagtc tatggcttgg tcagtacaga aggagtgggtg 360
tcctgctttg gagagaagac cccgcagcat cctggccctc ctgcttcatt gtccctgggc 420
aacaggtgcc acaacgttgt gacagctgta ggagcctggc cagctcatgg gagcatcctt 480
ggaaatgttc cagaagcccc tgtgggagct gatgtgttgg gggctggagg atgtgactgg 540
gcagacaaaag aggccttggc ccctgggcaa agggcaaagg tgcacattct tcttgagagt 600
tctggacagt ctgatccatc ctatgctgtc cttcctgaca gctgggcagc cacggagggtg 660
ttcccaactt acagatctca ggtctcctct ccccgcatcc cgggtagttc catctgggtt 720
ggcagtggtg ctggttggcc tatacttggg gaactcaggg aatgtgacca gatgttctcc 780
tgcatgttgc ccactgggtg tgcctccttc caggatccag gacgttatgg tgattatgac 840
ctccctatgg atgaggatga ggacatgacc aagaccggga ccttcttcgc agccaagatc 900
gtcattggca ttgcactggc aggcacatg ctggtctgcg gcatcggtaa ctttgtcttt 960
atcgctgccc tcaccgcta taagaagtgt cgcaacctca ccaatctgct cattgccaac 1020
ctggccatct ccgacttctt ggtggccatc atctgtctgc ctttcgagat ggactactac 1080
gtggtacggc agctctcctg ggagcatggc cacgtgctct gtgcctccgt caactacctg 1140
cgcaccgtct ccctctacgt ctccaccaat gccttgtgtg ccattgcat tgacaggtat 1200
ctcgccatcg ttcaccctt gaaaccacgg atgaattatc aaacggcctc cttcctgac 1260

```

```

gccttggtct ggatggtgct cattctcatt gccatcccat cggcttactt tgcaacagaa 1320
acggtccctct ttattgtcaa gagccaggag aagatcttct gtggccagat ctggcctgtg 1380
gatcagcagc tctactaaa gtcctacttc ctcttcatct ttggtgtcga gttcgtgggc 1440
cctgtggtca ccatgacctt gtgctatgcc aggatctccc gggagctctg gttcaaggca 1500
gtccctgggt tccagacgga gcagattcgc aagcggctgc gctgcccgag gaagacggtc 1560
ctggtgctca tgtgcattct cacggcctat gtgctgtgct gggcaccctt ctacggtttc 1620
accatcgctt gtgacttctt cccactgtg ttctgtgaagg aaaagcacta cctcactgcc 1680
ttctacgtgg tgcagtgcac cgccatgagc aacagcatga tcaacaccgt gtgcttctgt 1740
acggtcaaga acaacacccat gaagtacttc aagaagatga tgctgctgca ctggcgtccc 1800
tcccagcggg ggagcaagtc cagtgtgac cttgacctca gaaccaacgg ggtgcccacc 1860
acagaagagg tggactgtat caggctgaag tgaccactg gtgtcacaca attgaaaacc 1920
ccagtcagtc actcagagca tcaccacca tcaaccaagt tcataggctg catgggaaat 1980
gacatctgtg ttcatgcctc cccgtgccc tcaagaagcc gaatgctgca aagtctgaac 2040
atacaatgag actagacatg aaccaaatac gctgacattt actgatatcc gctcgacacc 2100
tactgtgtcc acaatcccaa caaggagatt agacacaagg agcagcaact gacatggact 2160
gaacatgtac tgtgtgcaag ccaaaccaat gagattaaca gggacagcag gagctgaatt 2220
atcttactat gtatcaaac tggtgttcac aaattaaact acagtcacac ttgggtcaca 2280
tcgttttatt tccatttcac tttt 2304

```

<210> 32

<211> 2322

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7482007CB1

<400> 32

```

cccatttcaa aaatggagaa gacagatcac tgccactgac caggaccgtg ggaggtgcc 60
cgtgatggtg aggcacatg ctaggagct gagctctgac ctctctgctg ggtgattctc 120
cacctctggg ctgctagatc tacttctctg atgccgtgaa gatcctcatg tatgaaaatg 180
aagtcccagg caaccatgat ttgctgctta gtgttctttc tgtccacaga atgttccac 240
tatagatcca agattcacct aaaagctgga gataaacttc aaagccctga agggaaaccc 300
aagactggaa ggatccaaga gaaatgcgaa ggaccttgta tttcttcttc caactgcagc 360
cagccctgtg ctaaggactt tcatggagaa ataggattta catgtaatca aaaaaagtgg 420
caaaaatcag ctgaaacatg tacaagcctt tctgtgaaa aactctttaa ggactcaact 480
ggtgcattct gcctttctgt agcagacca tctatacttc tgcatattct agactttcga 540
gctccagaga ccattgagag tgtagctcaa ggaatccgta agaactgcc ctttgattat 600
gcctgcatca ctgacatggt gaaatcatca gaaacaacat ctggaaatat tgcatttata 660
gtggagtatt taaaaatat ttctacagac ttgtctgata atgttactcg agagaaaatg 720
aagagctata gtgaagtggc caaccacatc ctgcacacag cagccatttc aaactgggct 780
ttcattccca acaaaaatgc cagctcggat ttgttgagc agtgaaattt gtttgccaga 840
caactccaca tccaataa ttctgagaac attgtgaatg aactcttcat tcagacaaaa 900
gggtttcaca tcaaccataa tacctcagag aaaagcctca atttctccat gagcatgaac 960
aataccacag aagatatctt agaatggta cagattccca ggcaagagct aaggaagctg 1020
tggccaaatg catcccaagc cattagcata gctttcccaa ccttgggggc tatcctgaga 1080
gaagccact tgcaaaatgt gagtcttccc agacaggtaa atggctcgtt gctatcagtg 1140
gttttaccag aaaggttgca agaaatcata ctacacttcg aaaagatcaa taaaaccgcg 1200
aatgccagag ccagtggtg ttggtggcac tccaagaaaa ggagatggga tgagaaagcg 1260
tgccaaatga tgttggtat caggaacgaa gtgaaatgcc gctgtaacta caccagtgtg 1320
gtgatgtctt ttccattct catgtcctcc aaatcgatga ccgacaaagt tctggactac 1380
atcacctgca ttgggtcag cgtctcaatc ctaagcttgg ttctttgect gatcattgaa 1440
gccacagtgt ggtcccgggt ggttgtgacg gagatatcat acatgcgtca cgtgtgcatc 1500
gtgaatatag cagtgtccct tctgactgcc aatgtgtggt ttatcatagg ctctcacttt 1560
aacattaagg cccaggacta caacatgtgt gttgcagtga catttttcag ccaacttttc 1620
tacctctctc tgtttttctg gattctcttc aaagcattgc tcatcattta tggaaatatt 1680
gtcattttcc ttaggatgat gaagtcccga atgatggta ttggccttgc cattggctat 1740
gggtgcccac tgatcatgct tgtcactaca gttgctatca cagggccagt gaaaggctac 1800
atgagacctg aggcctgttg gcttaactgg gacaatacca aagccctttt agcatttgcc 1860
atcccggcgt tcgtcattgt ggctgtaaat ctgattgtgg ttttggttgt tgctgtcaac 1920
actcagaggc cctctattgg cagttccaag tctcaggatg tggtcataat tatgaggatc 1980
agcaaaaatg ttgccatcct cactccactg ctgggactga cctggggttt tggaaatagcc 2040

```

```

actctcatag aaggcaattc cttgacgttc catataatth ttgccttgct caatgctttc 2100
cagggttttt tcatcctgct gtttggaaac attatggatc acaagataag agatgctttg 2160
aggatgagga tgtcttcact gaaggggaaa tcgagggcag ctgagaatgc atcactaggc 2220
ccaaccaatg gatctaaatt aatgaatcgt caaggatgaa atgctgcccc atttctcatg 2280
gatgtcctga gaccaagagg ggagatccag gagaaagagg cc 2322

```

```

<210> 33
<211> 2366
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 6769042CB1

```

```

<400> 33
atttaggtga cactatagaa gagcccagtg tgctggaaag gagatcgcca tgtacttcac 60
tgctgccatt ggaaagcatg ctttattgtc ttcaacgctg ccaagcctct tcatgacatc 120
cacagcaagc cccgtgatgc ccacagatgc ctaccatccc atcataacca acctgacaga 180
agagagaaaa accttccaaa gtcccggagt gatactgagt tacctccaaa atgtatccct 240
cagcttaccg agtaagtccc tctcggagca gacagccttg aatctcacca agaccttctt 300
aaaagccgtg ggagagatcc ttctactgcc tgggttgatt gctctgtcag aggacagcgc 360
cgtgggtactg agtctcatcg acaactattga caccgtcatg ggccatgtat cctccaacct 420
gcacggcagc acgccccagg tcaccgtgga gggctcctct gccatggcag agttttccgt 480
ggccaaaatc ctgcccaga ccggtgaattc ctcccattac cgcttcccgg cccacgggca 540
gagcttcatc cagatcccc acgaggcctt ccacaggcac gcctggagca ccgtcgtggg 600
tctgctgtac cacagcatgc actactacct gaacaacatc tggcccggcc acaccaagat 660
cgcgagggcc atgcatcacc aggactgcct gctgttcgcc accagccacc tgatttccct 720
ggaggtgtcc ccaccacca ccctgtctca gaacctgtcg ggctctccac tcattacggg 780
ccacctcaag cacagattga cacgtaagca gcacagttag gccaccaaca gcagcaaccg 840
agtcttcgtg tactgcgcct tccctggactt cagctccgga gaagggtctt ggtcgaacca 900
cggtctgtcg ctcacgagag gaaacctcac ctactccgtc tgccgctgca ctacacctac 960
caactttgcc atcctcatgc aggtgggtccc gctggagctt gcacgcggac accaggtggc 1020
gctgtcgtct atcagctatg tgggctgtct cctctccgtg ctctgcctgg tggccacgct 1080
ggtcaccttc gccgtgctgt cctccgtgag caccatccgg aaccagcgct accacatcca 1140
cgccaacctg tccttcgccg tgctgggtggc ccaggtcctg ctgctcatta gtttccgctt 1200
cgagccaggc acgaccccct gccaaagtga ggccgtgctc ctacactact tcttcttag 1260
tgcttcgca tggatgctgg tggaggggct gcacctctac agcatggtga tcaaggtctt 1320
tgggtcggag gacagcaagc accgttacta ctatgggatg ggatgggggt ttctctctt 1380
gatctgcac atttcactgt catttgccat ggacagttac ggaacaagca acaattgctg 1440
gctgtcgttg gcgagtggcg ccactctggc cttttagacc cctgccctgt ttgtcatcgt 1500
ggtcaacatt ggcatcctca tcgtgtgac cagagtcac tcacagatca gcgccgacaa 1560
ctacaagatc catggagacc ccagtgcctt caagttgacg gccaaaggcag tggccgtgct 1620
gctgcccatc ctgggtacct cgtgggtctt tggcgtgctt gctgtcaacg gttgtgctgt 1680
ggttttccag tacatgtttg ccacgtcaa ctccctgcag ggactgttca tattcctctt 1740
tcattgtctc ctgaattcag aggtgagagc cgccttcaag cacaaaatca aggtctggtc 1800
gtcacgagc agctccgccc gcacctcaa cgcgaagccc ttccactcgg acctcatgaa 1860
tgggacccgg ccaggcatgg cctccaccaa gctcagccct tgggacaaga gcagccactc 1920
tgccaccgc gtcgacctgt cagccgtgtg agccgggagg ctgccaacca ggccaggctg 1980
cgctcagaac acaccccccc aaacagaatg aaatgcccc cctttgcccc tggacctctt 2040
ccttgctgct gtctggacat ggggtgttgt gccccgagac agctgtcctc ccctgtgact 2100
ctggctgtgc gagcacactg ctcagcccag cagcctgatg cccaggccag cgtgggccct 2160
cctgccttgc atccacccgt gggctgagtg acttccctcg gggattccca ggacacagtg 2220
gcctgacttg tgatggtgcc cttgagcctc ccttcatcac tcagcatcag accagcgagg 2280
cagggcatcg gggccggtcc cgcagcccgg agggatgtca gctctgtgct ggggggttgg 2340
ggcccgcccc aagtgtcagg ccccg 2366

```

```

<210> 34
<211> 1458
<212> DNA
<213> Homo sapiens

```

```

<220>

```

<221> misc_feature

<223> Incyte ID No: 7476053CB1

<400> 34

```
atggaggccg ctagcctttc agtggccacc gccggcggtt cccttgccct gggaccgag 60
accagcagcg ggacccaag cccgagaggg atactcggtt cgaccccgag cggcgccgtc 120
ctgccggggc gagggccgcc cttctctgtc ttcacgggtc tgggtggtgac gctgctagtg 180
ctgctgatcg ctgccacttt cctgtggaac ctgctgggtc cggtcaccat cccgcggtc 240
cgtgccttcc accgcgtgcc gcataacttg gtggcctcga cggcctctc ggacgaacta 300
gtggcagcgc tggcgatgcc accgagcctg gcgagtgagc tgtcgaccgg gcgacgtcgg 360
ctgctggggc ggagcctgtg ccacgtgtgg atctccttcg acgcccgtg ctgccccgcc 420
ggcctcggga acgtggcggc catcgccctg ggccgcgacg gggccatcac acggcacctg 480
cagcacacgc tgcgcacccg cagccgcgcc tcgttgctca tgatcgcgct cgcccggtg 540
ccgtcggcgc tcatcgccct cgcgcgctg ctctttggcc gggcgaggt gtgcgacgt 600
cggctccagc gctgcagggt gagccgggaa ccctcctatg ccgccttctc caccgcggc 660
gccttcacc tgcgccttgg cgtggtgccg tttgtctacc ggaagatcta cgaggcgcc 720
aagtttcgtt tcggccgccc cggagagct gtgctgccg tgcggccac catgcagggt 780
aaggaaagcac ctgatgagc tgaagtgggt ttcacggcac attgcaaagc aacggtgtcc 840
ttccagtgga gcggggactc ctggcgggag cagaaggaga ggcgagcagc catgatggtg 900
ggaattctga ttggcgtggt tgtgctgtgc tggatccctc tottcctgac ggaactcatc 960
agccactct gtgcctgcag cctgcccccc atctggaaaa gcatatttct gtggcttggc 1020
tactccaatt ctttcttcaa cccctgatt tacacagctt ttaacaagaa ctacaacaat 1080
gccttcaaga gcctctttac taagcagaga tgaacacagg ggtagagag acatgggtag 1140
attttaagga ggaaggaact tggacttttt cgtcagtgat ctgagattct tccctccaca 1200
gctgagtgct aatgtgtgat tgagagttat accattgggc ctggactgta gaagcagcag 1260
agccaaggtt ctcaagaaag acagcaaagg tctggcagat gttgtaacta tgccttcttc 1320
ccatgtgcat ggcagacatt gccaatgggt catggcttgg ctcccactg agcaggaaact 1380
tggctcaga atcctttcca ggacagcacc ctaggcagct actgttgatt atttaaaatt 1440
gatgcaagac ttgaaaaa 1458
```

<210> 35

<211> 975

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7480410CB1

<400> 35

```
atgggcatgg agggctcttct ccagaactcc actaacttcg tcttcacagg cctcatcacc 60
catcctgcct tccccgggct tctctttgca atagtcttct ccatctttgt ggtggctata 120
acagccaact tggctcatgat tctgctcatt cacatggact cccgcctcca cactcccatg 180
tacttcttgc tcagccagct ctccatcatg gataccatct acatctgtat cactgtcccc 240
aagatgctcc aggacctcct gtccaaggac aagaccattt ccttcctggg ctgtgcagtt 300
cagatcttcc tctacctgac cctgattgga ggggaattct tctgctggg tctcatggcc 360
tatgaccgct atgtggctgt gtgcaaccct ctacggtagc ctctcctcat gaaccgagg 420
gtttgcttat tcatggtggg cggtcctgg gttaggtggt ccttgatggg gttcatgctg 480
actcctgtca ctatgagttt ccccttctgt agatcccag agatcaatca ctttttctgt 540
gagatccag ccgtgctgaa gttgtcttgc acagacagct cactctatga gacctgatg 600
tatgctgct gcgtgctgat gctgcttacc cctctatctg tcatctctgt ctctacacg 660
cacatcctcc tgactgtcca caggatgaac tctgctgagg gccggcgcaa agccttggct 720
acgtgttccct cccacattat ggtggtgagc gttttctacg gggcagcctt ctacaccaac 780
gtgctgcccc actcctacca cactccagag aaagataaag tgggtgtctgc cttctacacc 840
atcctcacc ccatgctcaa cccactcatc tacagcttga ggaataaaga tgtggctgca 900
gctctgagga aagtactagg gagatgtggg tctcccaga gcatcagggt ggcgactgtg 960
atcaggaagg gctag 975
```

<210> 36

<211> 948

<212> DNA

<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 55036418CB1

<400> 36
atggagacgt ggggtgaacca gtcctacaca gatggcttct tcctcttagg catcttctcc 60
cacagtactg ctgaccttgt cctcttctcc gtggttatgg cggctcttcac agtggccctc 120
tgtgggaatg tcctcctcat ctctcctatc tacatggacc ctcaccttca caccctcatg 180
tacttcttcc tcagccagct ctccctcatg gacctcatgt tggctctgtac caatgtgcc 240
aagatggcag ccaacttcct gtctggcagg aagtccatct cctttgtggg ctgtggcata 300
caaattggcc tctttgtctg tcttgtggga tctgaggggc tcttgtctgg actcatggct 360
tatgaccgct atgtggccat tagccacca cttcactatc ccatcctcat gaatcagagg 420
gtctgtctcc agattactgg gagctcctgg gcctttggga taatcgatgg cttgatccag 480
atggtggttag taatgaattt cccctactgt ggcttgagga aggtgaacca tttcttctgt 540
gagatgctat ccttggttgaa gctggcctgt gtagacacat ccctgtttga gaagggtgata 600
tttgccttct gtgtcttcat gcttctcttc ccattctcca tcactcgtggc ctccatgtct 660
cacattctag ggactgtgct gcaaatgcac tctgctcagg cctggaaaaa ggccctggcc 720
acctgtcctc cccacctgac agctgtcacc ctcttctatg gggcagccat gttcatctac 780
ctgaggccta ggcactaccg ggccccagc catgacaagg tggcctctat cttctacacg 840
gtccttactc ccatgctcaa cccctcatt tacagcttga ggaacaggga ggtgatgggg 900
gcactgagga aggggctgga ccgctgcagg atcggcagcc agcactga 948

<210> 37
<211> 1086
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 7481701CB1

<400> 37
ggctctattc agacgetggc ttcttgtaag tgtattcctt tatccaatag tagatgcctc 60
ctagaaggct tgagtgcact ggaaattgaa actctcactt tcaacttgga gatggagagc 120
cccaatcaaa ccaccattca ggagtttata ttctccgctt tcccttattc ctgggttaag 180
tctgttgtct gctttgttcc actgctcttc atctatgctt tcattgttgt tggaaacctg 240
gtcatcatca cagtggcca gttgaatact caccctccaca ctcccatgta tacttttata 300
agtgtctctt ctttctgga gatttgggtat accacagcca caatcccaaa gatgctgtct 360
agcctgttta gtgagaggag catttccttc aatggttgct tcctgcagat gtatttcttc 420
cattccaccg gcatctgtga ggtgtgtctc ttgacagtta tggcctttga ccaactacctg 480
gccatattgca gccctcttca ttatccctct atcatgaccc ccaagctatg tacccaactg 540
actttaagtt gctgtgtttg tggctttatc acaccgttc ctgagattgc ctggatctct 600
acactgccat tttgtgggtc gaatcacctt gaacatatct tctgtgactt cctccagtg 660
ctgcgtctgg cctgcacaga cacacgagcc atcgtcatga ttcaggtagt ggatgtcatt 720
catgcagtgg agattattac agctgtgatg ctcatcttca tgtcctacga tggattattg 780
gctgtaattc tacgtattca ttcagctgga ggccgccgca cagcattttc cacgtgtgtc 840
tctcacttca ttgtcttttc gctcttctt ggagctgtga ctctcatgta cctacgcttc 900
tctgccacct actctttgt ctgggata gccattgctc tggcctttgc agttttgtct 960
cccttcttca acccattat ctatagcctg aggaataaag aaataaaaga agctataaaa 1020
aagcacatag gtcaagctaa gatatttttt tccgtaagac caggacctc aagtaagata 1080
ttttag 1086

<210> 38
<211> 1529
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 7481774CB1

<400> 38
aaggagacc acagtgagag ggagccctga gcagaagtaa ggctgtcaca aggctggaag 60

cagagaacat	ccccatggaa	ctgaagacag	catgctgcat	ccctgggagg	agggagctct	120
taaggaagt	ccaaggattg	atatttctgt	tcagctgcag	tagagatgga	tggaaccatg	180
gcagcacc	acccatttca	tcctactggg	attctctgac	cgaccccatc	tggaaggat	240
cctctttgtg	gtcatcctga	tcgcgtacct	cctgaccctc	gtaggcaaca	ccaccatcat	300
cctggtgtcc	cggctggacc	cccacctcca	caccccatg	tactttctcc	tcgcccacct	360
ttccttcctg	gacctcagtt	tcaccaccag	ctccatcccc	cagctgctct	acaaccttaa	420
tggtgtgac	aagaccatca	gctacatggg	ctgtgccatc	cagctcttcc	tggtcctggg	480
tctgggtggt	gtggagtgcc	tgcttctggc	tgtcatggcc	tatgaccggt	gtgtggctat	540
ctgcaagccc	ctgcactaca	tggtgatcat	gaaccccagg	ctctgccggg	gcttggtgtc	600
agtgacctgg	ggctgtgggg	tggccaactc	cttggccatg	tctcctgtga	ccctgcgctt	660
accccgctgt	gggcaccacg	aggtggacca	cttcctgtgt	gagatgcccg	ccctgatccg	720
gatggcctgc	atcagcactg	tggccatcga	cggcaccgtc	tttgtcctgg	cgggtgggtgt	780
tgtgctgtcc	cccttggtgt	ttatcctgct	ctcttacagc	tacattgtga	gggctgtgtt	840
acaaattcgg	tcagcatcag	gaaggcagaa	ggccttcggc	acctgcgggt	cccattctcac	900
tgtggtctcc	cttttctatg	gaaacatcat	ctacatgtac	atgcagccag	gagccagttc	960
ttcccaggac	cagggcaagt	tcctcacgct	cttctacaac	attgtcaccc	ccctcctcaa	1020
tcctctcatc	tacaccctca	gaaacagaga	ggtgaagggg	gcactgggaa	ggttgcttct	1080
ggggaagaga	gagctaggaa	aggagttaaag	gcattctccac	ctgacttcac	ctccatccag	1140
ggccactggc	agcatctgga	acggctgaat	tccagctgat	attagccac	gactcccaac	1200
ttgccttttt	ctggactttt	gtgaggctgt	ttcagttctg	acattatgtg	tttttgtgtg	1260
tgtctttaa	attgagacgg	ggtctcactc	tgtcacctag	ggtggagtgc	agtggtgcca	1320
ccatagctcc	ttcgactatt	gggcttaagc	gatcctcccc	cacctcagcc	ttccaagtaa	1380
ctgggactac	aggtgtgcat	cactggcagt	gggaattgtg	gcttttctgt	cttctatgga	1440
gacggggtct	tgctgtgttg	accaggctgg	tcccâaactc	ctggcctcat	gtgatcctcc	1500
tgccatggcc	tcctaaagtt	ctgggatta				1529